Attorney Docket No.: 2307o-099910US

PATENT APPLICATION NUCLEIC ACIDS THAT CONTROL PLANT DEVELOPMENT

Inventor(s):

Robert L. Fischer, a citizen of the United States residing at: 1423 Scott Street, El Cerrito, CA 94530

Yeonhee Choi, a citizen of South Korea residing at: 4 Anchor Dr. #442, Emeryville, CA 94608

Mike Hannon, a citizen of the United States residing at: 309 Pearl Drive, Livermore, CA 94550

Jack Kishiro Okamuro, a citizen of the United States residing at: 307 Ocho Rios Way, Oak Park, CA 91377

Tatiana Valerievna Tatarinova, a citizen of Russia residing at 836 North Beaudry Avenue Apt. 2, Los Angeles, CA 90012

Assignee:

THE REGENTS OF THE UNIVERSITY OF CALIFORNIA 1111 Franklin Street, Fifth Floor Oakland, CA 94607-5200

Entity:

TOWNSEND and TOWNSEND and CREW LLP Two Embarcadero Center, 8th Floor San Francisco, California 94111-3834 Tel: 415-576-0200

NUCLEIC ACIDS THAT CONTROL PLANT DEVELOPMENT

CROSS-REFERENCES TO RELATED APPLICATIONS

1266-

This application is a continuation-in-part of U.S. Patent Application No. 09/553,690, filed April 21, 2000, the contents of which are incorporated by reference.

STATEMENT AS TO RIGHTS TO INVENTIONS MADE UNDER FEDERALLY SPONSORED RESEARCH AND DEVELOPMENT

10

5

This invention was made with Government support under Grant No. 97-35304-4941, awarded by the United States Department of Agriculture. The government has certain rights in this invention.

FIELD OF THE INVENTION

15

This invention is directed to plant genetic engineering. It relates to, for example, modulating seed (and in particular endosperm, embryo and seed coat) development, flowering time, chromosomal DNA methylation and modulating transcription in plants.

BACKGROUND OF THE INVENTION

20

25

30

A fundamental problem in biology is to understand how seed development. In flowering plants, the ovule generates the female gametophyte, which is composed of egg, central, synergid and antipodal cells (Reiser, et al., Plant Cell, 1291-1301 (1993)). All are haploid except the central cell which contains two daughter nuclei that fuse prior to fertilization. One sperm nucleus fertilizes the egg to form the zygote, whereas another sperm nucleus fuses with the diploid central cell nucleus to form the triploid endosperm nucleus (van Went, et al., Embryology of Angiosperms, pp. 273-318 (1984)). The two fertilization products undergo distinct patterns of development. In Arabidopsis, the embryo passes through a series of stages that have been defined morphologically as preglobular, globular, heart, cotyledon and maturation (Goldberg, R. B., et al., Science (1994) 266: 605-614; Mansfield, S. G., et al., Arabidopsis: An Atlas of Morphology and Development, pp. 367-383 (1994)). The primary endosperm nucleus undergoes a series of mitotic divisions to produce nuclei that migrate into the expanding central cell (Mansfield, S. G., et al., Arab Inf Serv 27: 53-64 (1990); Webb, M. C., et al., Planta 184:187-195 (1991)). Cytokinesis sequesters endosperm cytoplasm and nuclei

into discrete cells (Mansfield, S. G., et al., Arab Inf Serv 27:65-72 (1990)) that produce storage proteins, starch, and lipids which support embryo growth (Lopes, M. A. et al., Plant Cell 5:1383-1399 (1993)). Fertilization also activates development of the integument cell layers of the ovule that become the seed coat, and induces the ovary to grow and form the fruit, or silique, in Arabidopsis.

Of particular interest are recent discoveries of genes that control seed, and in particular endosperm, development. For instance, *MEDEA* (*MEA*) (also known as *FIE1* (see, e.g., copending U.S. patent application 09/071,838) and F644 (see, e.g., Kiyosue T, *et al.* (1999) *Proc Natl Acad Sci U S A* 96(7):4186-91) encodes an Arabidopsis SET domain polycomb protein that appears to play a role in endosperm development. Inheritance of a maternal loss-of-function mea allele results in embryo abortion and prolonged endosperm production, irrespective of the genotype of the paternal allele. Thus, only the maternal wild-type MEA allele is required for proper embryo, endosperm, and seed coat development (Kinoshita T, *et al.* (1999) *Plant Cell* 10:1945-52). These results reveal functions for plant polycomb proteins in the suppression of central cell proliferation and endosperm development (Kiyosue T, *et al. supra*).

Another gene product that controls seed development is FIE, also known as FIE3 (see, e.g., copending U.S. patent application 09/071,838). The FIE protein is a homolog of the WD motif-containing Polycomb proteins from Drosophila and mammals (Ohad, N. et al. Plant Cell 11(3):407-16 (1999)). In Drosophila, these proteins function as repressors of homeotic genes. Loss of function mutations in the FIE gene result in endosperm phenotypes that are identical to medea loss of function mutations. A female gametophyte with a loss-of-function allele of fie undergoes replication of the central cell nucleus and initiates endosperm development without fertilization. These results suggest that the FIE Polycomb protein functions to suppress a critical aspect of early plant reproduction, namely, endosperm development, until fertilization occurs. Moreover, hypomethylation of fie mutants leads to the development of differentiated endosperm. Vinkenoog et al., Plant Cell 12:2271-2282 (2000).

Control of the expression of genes that control egg and central cell differentiation, or those that control reproductive development, i.e. embryo, endosperm and seed coat, is useful in the production of plants with a range of desired traits. These and other advantages are provided by the present application.

SUMMARY OF THE INVENTION

This invention provides isolated nucleic acids comprising a polynucleotide sequence, or its complement, encoding a DMT polypeptide comprising an amino acid sequence with at least 70% sequence identity to at least one of the following consensus sequences:

DMT Domain A

KV<1>(I,I)D(D,p) (E,V)T<3>W<1>(L,V)L(M,1) (E,d)<0-2>D(K,e)<1>(K,t)<1>(K,a) (W,k) (W,l)<1>(E,k)ER<2>F<1>(G,t)R<1>(D,n) (S,l)FI(A,n)RM(H,r)<1>(V,l)QG(D,n)R<1>F<1>(P,q)WKGSVVDSV(I,V)GVFLTQN(V,t)D(H,y) (L,s)SS(S,n)A(F,y)M<1>(L,V)A(A,s)<1>FP

DMT Domain B

W(D,n)<1>(L,f)R<5>E<3-

6>D(S,t)<1>(D,n) (Y,w)<3>R<10>I<2>RG(M,q) (N,f)<2>L(A,s)<1>RI<2-12>FL<3>V<2>(H,n)G<1>IDLEWLR<2>(P,d) (P,s) (D,h)<1>(A,v)K<1>(Y,f)LL(S,e) (I,f)<1>G(L,i)GLKS (V,a)ECVRLL<1>L(H,k)<2>AFPVDTNVGRI (A,c)VR(M,1)G(W,1)VPL(Q,e)PLP<2>(L,v)Q(L,m)H(L,q)L(E,f)<1>YP<1>(L,m) (E,d) (S,n) (I,v)QK(F,y)LWPRLCKL(D,p)Q<1>TLYELHY (Q,h) (L,m)ITFGK<0-2>FCTK<2>PNCNACPM(R,k)<0-2>EC(R,k) (H,y) (F,y) (A,s)SA<1>(A,v)<0-10>S(A,s) (R,k)<1>(A,1)L(P,e)<1>(P,t) (N,f)<1>(N,f)<1>(N,f)<1>(N,f)<1>(N,f)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1>(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g)<1<(N,g

20

15

5

10

DMT Domain C.

P(I,1)(I,v)E(E,f)P<1>(S,t)P<2-5>E<0-15>(D,a)IE(D,e)<423>(I,v)P<1>I<1>(L,f)(N,d)<8-17>(S,a)<1>(A,d)LV<8>(I,1)P<25>(K,r)(L,m)K<4>LRTEH<1>V(Y,f)(E,v)LPD<1>H<1>(L,i)L(E,k)<1>(D,e)D(P,i)<2>YLL(A,s)
IW(T,q)P(G,d)(E,g)<6-8>(P,s)<3>C<610>(M,1)C<4>C<2>C<3>(R,k)E<5>(V,f)RGT(L,i)L<022>(L,v)FADH<1>(S,t)(S,r)<2>PI<3>(R,t)<3>(W,k)<1>L<1>(R,k)R<4>G(T,s)(S,t)<2>(S,t)
I(F,c)(R,k)(G,1)L<1>(T,v)<2>I<2>(C,n)F(W,q)<1>G(F,y)(V,1)C(V,1)R<1>F(E,d)<3>(R,t)<3>(R,t)<3>(R,t)<3<(R,t)<3<(R,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,t)<4<(T,s)(S,

30

35

25

In some embodiments, the nucleic acids of the invention do not encode a polypeptide at least 40% identical to SEQ ID NO:2, or alternatively at least 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 99% or 100% to SEQ ID NO:2. In some embodiments, the DMT polypeptide comprises an amino acid sequence 100% identical to the above-listed consensus sequences.

In some embodiments, the DMT polypeptides ar at least 45%,50%,55%,60%,65%,70%,75%,80%,85%,90%,95%,97%,98%,99% or 100% identical to DMT domains A, B and/or C.

10

15

20

25

30

number;

In one aspect, the invention provides DMt polypeptides capable of exhibiting at least one of the following biological activities: (a) glycosylase activity; (b) demethylation of polynucleotides; (c) DNA repair; (d) wherein expression of the polypeptide in a plant modulates organ identity; (e) wherein expression of the polypeptide in a plant modulates organ number; (f) wherein expression of the polypeptide in a plant modulate meristem stem and/or activity: (g) wherein enhanced expression of the polypeptide in a plant results in a delay in flowering time; (h) wherein introduction of the polypeptide into a cell results in modulation of methylation of chromosomal DNA in the cell; (i) wherein reduction of expression of the polypeptide in a plant results in modulation of endosperm development; (i) wherein expression of the polypeptide in an Arabidopsis leaf results in modulation of expression of the MEDEA gene. In some aspects, the polypeptide comprises either a (i) basic region; (ii) nuclear localization signal; (iii) leucine zipper; (iv) helix-hairpin-helix structure; (v) glycine-proline rich loop with a terminal aspartic acid or (vi) helix that is capable of binding DNA. In one aspect, the invention provides methods of modulating in a plant one or more of the following: (a) DNA repair; (b) wherein expression of the polypeptide in a plant modulates organ identity;

(c) wherein expression of the polypeptide in a plant modulates organ

10

15

20

25

30

- (d) wherein expression of the polypeptide in a plant modulate meristem stem and/or activity;
- (e) wherein enhanced expression of the polypeptide in a plant results in a delay in flowering time;
- (f) wherein introduction of the polypeptide into a cell results in modulation of methylation of chromosomal DNA in the cell;
- (g) wherein reduction of expression of the polypeptide in a plant results in modulation of endosperm development;
- (h) wherein expression of the polypeptide in an Arabidopsis leaf results in expression of the MEDEA gene,

wherein the method comprises:,

- (a) introducing into a plant cell a nucleic acid of claim 1; and
- (b) generating conditions where the plant cell can transcribe the nucleic acid described above.

In some embodiments, the polypeptides comprise between 1500 and 2000 amino acids. In some aspects, the polypeptide has glycosylase activity. In some embodiments, introduction of the nucleic acid into a cell results in modulation of methylation of chromosomal DNA in the cell. In some embodiments, enhanced expression of the nucleic acids of the invention into a plant results in a delay in flowering time. In some embodiments, reduction of expression of a DMT polypeptide in a plant results in enhanced endosperm development. In addition, in some embodiments, expression of the nucleic acid of the invention in an *Arabidopsis* leaf results in expression of the *MEDEA* gene.

This invention provides isolated nucleic acids comprising a polynucleotide sequence, or its complement, encoding a DMT polypeptide exhibiting at least 60% sequence identity to SEQ ID NO:2 or exhibiting at least 70% sequence identity to at least one of DMT domain A, B, or C. For instance, the nucleic acid can encode the DMT polypeptide displayed in SEQ ID NO:2. In one aspect, the polynucleotide sequence comprises SEQ ID NO:5 or SEQ ID NO:1. In some aspects of the invention, the nucleic acid further comprises a promoter operably linked to the polynucleotide. In some embodiments, the promoter is constitutive. In other embodiments, the promoter is from a DMT gene. For example, the promoter can comprise a polynucleotide at least 70% identical to SEQ ID NO:3. In some aspects, the promoter comprises SEQ ID NO:3. In some aspects of this invention, the promoter further comprises a polynucleotide at least

10

15

20

25

30

70% identical to SEQ ID NO:4. For example, in some aspects the promoter comprises SEQ ID NO:4. In some aspects, the polynucleotide sequence is linked to the promoter in an antisense orientation.

The invention also provides an isolated nucleic acid molecule comprising a polynucleotide sequence exhibiting at least 60% sequence identity to SEQ ID NO:1.

The invention also provides an expression cassette comprising a promoter operably linked to a heterologous polynucleotide sequence, or complement thereof, encoding a DMT polypeptide exhibiting at least 60% sequence identity to SEQ ID NO:2. For instance, the nucleic acid can encode the DMT polypeptide displayed in SEQ ID NO:2. In some aspects, the polynucleotide sequence comprises SEQ ID NO:5 or SEQ ID NO:1. In some aspects of the invention, the nucleic acid further comprises a promoter operably linked to the polynucleotide. In some embodiments, the promoter is constitutive. In other embodiments, the promoter is from a DMT gene. For example, the promoter can comprise a polynucleotide at least 70% identical to SEQ ID NO:3. In some aspects, the promoter comprises SEQ ID NO:3. In some aspects of this invention, the promoter further comprises a polynucleotide at least 70% identical to SEQ ID NO:4. For example, in some aspects the promoter comprises SEQ ID NO:4. In some aspects, the polynucleotide sequence is linked to the promoter in an antisense orientation.

The invention also provides an expression cassette for the expression of a heterologous polynucleotide in a plant cell. In some aspects, the expression cassette comprises a promoter polynucleotide at least 70% identical to SEQ ID NO:3 that is operably linked to a heterologous polynucleotide. In some aspects, the promoter comprises SEQ ID NO:3. In some aspects, the promoter further comprises a polynucleotide at least 70% identical to SEQ ID NO:4. For instance, in some embodiments, the promoter comprises SEQ ID NO:4. In some aspects, the promoter further comprises a polynucleotide at least 70% identical to SEQ ID NO:6. In some aspects, the promoter comprises SEQ ID NO:6.

The present invention also provides a host cell comprising an exogenous polynucleotide sequence comprising a polynucleotide sequence, or complement thereof, encoding a DMT polypeptide exhibiting at least 60% sequence identity to SEQ ID NO:2 or exhibiting at least 70% sequence identity to at least one of DMT domain A, B, or C. In some aspects of the invention, the nucleic acid further comprises a promoter operably linked to the polynucleotide sequence. In some aspects, the promoter is constitutive. In some aspects, the promoter comprises a polynucleotide at least 70% identical to SEQ ID

10

15

20

25

30

NO:3. The promoter, for instance, can comprise SEQ ID NO:3. In some aspects, the promoter further comprises a polynucleotide at least 70% identical to SEQ ID NO:4. For instance, in some embodiments, the promoter comprises SEQ ID NO:4. In some aspects, the promoter is operably linked to the exogenous polynucleotide sequence in an antisense orientation.

The present invention also provides an isolated polypeptide comprising an amino acid sequence at least 60% identical to SEQ ID NO:2 or an amino acid sequence at least 70% sequence identical to at least one of DMT domain A, B, or C and capable of exhibiting at least one biological activity of the polypeptide displayed in SEQ ID NO:2, or fragment thereof. The present invention also provides for an antibody capable of binding such polypeptides.

The present invention also provides a method of introducing an isolated nucleic acid into a host cell comprising, (a) providing an isolated nucleic acid or its complement, encoding a DMT polypeptide exhibiting at least 60% sequence identity to SEQ ID NO:2 or exhibiting at least 70% sequence identity to at least one of DMT domain A, B, or C and (b) contacting the nucleic acid with the host cell under conditions that permit insertion of the nucleic acid into the host cell.

The present invention also provides a method of modulating transcription, comprising introducing into a host cell an expression cassette comprising a promoter operably linked to a heterologous DMT polynucleotide, the heterologous DMT polynucleotide encoding a DMT polypeptide at least 60% identical to SEQ ID NO:2 or at least 70% sequence identical to at least one of DMT domain A, B, or C, and detecting a host cell with modulated transcription. In some aspects of the invention, the heterologous DMT polynucleotide encodes SEQ ID NO:2. In some aspect, the polynucleotide sequence comprises SEQ ID NO:5 or SEQ ID NO:1. In some aspects, the expression cassette is introduced into a host cell by *Agrobacterium*. In some aspects, the expression cassette is introduced by a sexual cross. In some aspects of the method of the invention, modulating transcription results in the modulation of endosperm development in a plant. In some aspects, endosperm development is enhanced. In other aspects, endosperm development is decreased. In some aspects of the invention, the promoter is operably linked to the DMT polynucleotide in an antisense orientation.

The present invention also provides a method of detecting a nucleic acid in a sample, comprising (a) providing an isolated nucleic acid molecule comprising a polynucleotide sequence, or its complement, encoding a DMT polypeptide exhibiting at

10

15

20

25

30

least 60% sequence identity to SEQ ID NO:2 or exhibiting at least 70% sequence identity to at least one of DMT domain A, B, or C, (b) contacting the isolated nucleic acid molecule with a sample under conditions that permit a comparison of the sequence of the isolated nucleic acid molecule with the sequence of DNA in the sample, and (c) analyzing the result of the comparison. In some aspects of the method, the isolated nucleic acid molecule and the sample are contacted under conditions that permit the formation of a duplex between complementary nucleic acid sequences.

The present invention also provides a transgenic plant cell or transgenic plant comprising a polynucleotide sequence, or its complement, encoding a DMT polypeptide exhibiting at least 60% sequence identity to SEQ ID NO:2 or exhibiting at least 70% sequence identity to at least one of DMT domain A, B, or C. For instance, the nucleic acid can encode the DMT polypeptide displayed in SEQ ID NO:2. In one aspect, the polynucleotide sequence comprises SEQ ID NO:5 or SEQ ID NO:1. In some aspects of the invention, the nucleic acid further comprises a promoter operably linked to the polynucleotide. In some embodiments, the promoter is constitutive. In other embodiments, the promoter comprises a polynucleotide at least 70% identical to SEQ ID NO:3. In some aspects, the promoter comprises SEQ ID NO:3. In some aspects of this invention, the promoter further comprises a polynucleotide at least 70% identical to SEQ ID NO:4. For example, in some aspects the promoter comprises SEQ ID NO:4. In some aspects, the polynucleotide sequence is linked to the promoter in an antisense orientation. The present invention also provides a plant that is regenerated from a plant cell as described above.

The present invention also provides an expression cassette for the expression of a heterologous polynucleotide in a plant cell, wherein the expression cassette comprises a promoter at least 70% identical to SEQ ID NO:3 and the promoter is operably linked to a heterologous polynucleotide. In some embodiments, the promoter comprises a polynucleotide at least 70% identical to SEQ ID NO:4 and/or SEQ ID NO:6. In some embodiments, the promoter specifically directs expression of the heterologous polynucleotide in a female gametophyte when the expression cassette is introduced into a plant.

DEFINITIONS

The phrase "nucleic acid sequence" refers to a single or double-stranded polymer of deoxyribonucleotide or ribonucleotide bases read from the 5' to the 3' end. It

10 -

15

20

25

30

includes chromosomal DNA, self-replicating plasmids, infectious polymers of DNA or RNA and DNA or RNA that performs a primarily structural role.

A "promoter" is defined as an array of nucleic acid control sequences that direct transcription of an operably linked nucleic acid. As used herein, a "plant promoter" is a promoter that functions in plants. Promoters include necessary nucleic acid sequences near the start site of transcription, such as, in the case of a polymerase II type promoter, a TATA element. A promoter also optionally includes distal enhancer or repressor elements, which can be located as much as several thousand base pairs from the start site of transcription. A "constitutive" promoter is a promoter that is active under most environmental and developmental conditions. An "inducible" promoter is a promoter that is active under environmental or developmental regulation. The term "operably linked" refers to a functional linkage between a nucleic acid expression control sequence (such as a promoter, or array of transcription factor binding sites) and a second nucleic acid sequence, wherein the expression control sequence directs transcription of the nucleic acid corresponding to the second sequence.

The term "plant" includes whole plants, plant organs (e.g., leaves, stems, flowers, roots, etc.), seeds and plant cells and progeny of same. The class of plants which can be used in the method of the invention is generally as broad as the class of flowering plants amenable to transformation techniques, including angiosperms (monocotyledonous and dicotyledonous plants), as well as gymnosperms. It includes plants of a variety of ploidy levels, including polyploid, diploid, haploid and hemizygous.

A polynucleotide sequence is "heterologous to" an organism or a second polynucleotide sequence if it originates from a foreign species, or, if from the same species, is modified from its original form. For example, a promoter operably linked to a heterologous coding sequence refers to a coding sequence from a species different from that from which the promoter was derived, or, if from the same species, a coding sequence which is different from any naturally occurring allelic variants.

A polynucleotide "exogenous to" an individual plant is a polynucleotide which is introduced into the plant, or a predecessor generation of the plant, by any means other than by a sexual cross. Examples of means by which this can be accomplished are described below, and include Agrobacterium-mediated transformation, biolistic methods, electroporation, in planta techniques, and the like. "Exogenous," as referred to within, is any polynucleotide, polypeptide or protein sequence, whether chimeric or not, that is initially or subsequently introduced into the genome of an individual host cell or the

10

15

20

25

30

organism regenerated from said host cell by any means other than by a sexual cross. Examples of means by which this can be accomplished are described below, and include Agrobacterium-mediated transformation (of dicots - e.g. Salomon et al. EMBO J. 3:141 (1984); Herrera-Estrella et al. EMBO J. 2:987 (1983); of monocots, representative papers are those by Escudero et al., Plant J. 10:355 (1996), Ishida et al., Nature Biotechnology 14:745 (1996), May et al., Bio/Technology 13:486 (1995)), biolistic methods (Armaleo et al., Current Genetics 17:97 1990)), electroporation, in planta techniques, and the like. Such a plant containing the exogenous nucleic acid is referred to here as a T0 for the primary transgenic plant and T1 for the first generation. The term "exogenous" as used herein is also intended to encompass inserting a naturally found element into a non-naturally found location.

The phrase "host cell" refers to a cell from any organism. Preferred host cells are derived from plants, bacteria, yeast, fungi, insects or other animals, including humans. Methods for introducing polynucleotide sequences into various types of host cells are well known in the art.

The "biological activity of a polypeptide" refers to any molecular activity or phenotype that is caused by the polypeptide. For example, the ability to transfer a phosphate to a substrate or the ability to bind a specific DNA sequence is a biological activity. One biological activity of DMT is glycosylase activity, i.e., cleavage of the nucleotide base from the nucleotide sugar). Another biological activity of DMT is to demethylate nucleotides (e.g., DMT has 5'-methylcytosine glycosylase activity). In addition, DMT has the ability to modulate endosperm production, as described herein, and to modulate flowering time in plants. For example, when DMT expression or DMT activity is increased in a plant, the flowering time of the plant is delayed. Moreover, expression of a DMT polypeptide in a plant tissue (e.g., a leaf) that does not typically express the *MEDEA* gene (Grossniklaus U, *et al.*, *Science* 280(5362):446-50 (1998)) results in the expression of *MEDEA*.

Additional biological activities of DMT polypeptides include: nuclear localization (e.g., as localized by amino acids 43-78 of SEQ ID NO:2); the ability to modulate plant organ size and/or number; the ability to modulate meristem size and/or activity; and to perform DNA repair, including nucleotide methylation or demethylation and/orrepair and/or removal of mis-matched nucleotides from DNA.

An "expression cassette" refers to a nucleic acid construct, which when introduced into a host cell, results in transcription and/or translation of an RNA or

10

15

20

30

35

polypeptide, respectively. Antisense or sense constructs that are not or cannot be translated are expressly included by this definition.

A "DMT nucleic acid" or "DMT polynucleotide sequence" of the invention is a subsequence or full length polynucleotide sequence of a gene which encodes a polypeptide involved in control of reproductive development and which, when the maternal allele is mutated by when DMT activity is reduced or eliminated in a maternal tissue or plant, allows for increased production of the endosperm and/or abortion of the embryo. In addition, overexpression of DMT in plants results in delayed time to flowering. Moreover, DMT is necessary and sufficient for expression of MEDEA in a plant cell. An exemplary nucleic acid of the invention is the Arabidopsis DMT sequence (SEO ID NO:1). Additional DMT nucleic acid sequences from a variety of plant species are also provided (e.g., SEQID NOs: 7-70). DMT polynucleotides are defined by their ability to hybridize under defined conditions to the exemplified nucleic acids or PCR products derived from them. A DMT polynucleotide is typically at least about 30-40 nucleotides to about 7000, usually less than about 10,000 nucleotides in length. More preferably, DMT polynucleotides contain a coding sequence of from about 100 to about 5500 nucleotides, often from about 500 to about 3600 nucleotides in length. A DMT polypeptide is typically at least 500 amino acids, typically at least 1000 amino acids, more typically at least 1500 amino acids. In some embodiments, a DMT polypeptide comprises fewer than 2000 amino acids, more typically fewer than 3000 amino acid and still more typically fewer than 5000 or 7500 amino acid in length.

As described below, *DMT* nucleic acid sequences encode polypeptides with substantial identity to at least one of following the consensus sequences:

DMT Domain A

DMT Domain B

W(D,n)<1>(L, t)R<5>E<3-6>D(S,t)<1>(D,n)(Y,w)<3>R<10>I<2>RG(M,q)(N,f)<2>L(A,s)<1>RI<2-12>FL<3>V<2>(H,n)G<1>IDLEWLR<2>(P,d)(P,s)(D,h)<1>(A,v)K<1>(Y,f)LL(S,e)(I,f)<1>G(L,i)GLKS(V,a)ECVRLL<1>L(H,k)<2>AFPVDTNVGRI(A,c)VR(M,1)G(W,1)VPL(Q,e)PLP<2>(L,v)Q (L,m)H(L,q)L(E,f)<1>YP<1>(L,m)(E,d)(S,n)(I,v)QK(F,y)LWPRLCKL(D,p)Q<1>TLYELHY(Q,h

) (L,m)ITFGK<0-2>FCTK<2>PNCNACPM(R,k)<0-2>EC(R,k)(H,y)(F,y)(A,s)SA<1>(A,v)<0-10>S(A,s)(R,k)<1>(A,1)L(P,e)<1>(P,t)

DMT Ddmain C.

In addition, the following cansensus sequence spanning all three domains

15 were identified:

<9-14>(T,q)(A,i)(S,k)(I,l)<3>(A,r)(S,k)(S,r)(P,k)<2>(K,f)<2>(E,l)K<0-1>K<03>(P,r)<2>(P,r)<1>(K,r)(K,r)(G,d)(R,k)<1>(G,v)<1>(K,g)<35>(P,s)(P,k)<3>(S,n)<1>(I,l)<0-2>(Q,d)<9>(P,q)<4>(K,a)(P,s)<14-16>(P,a)<4>L<010>D<1>(I,l)<0-4>(L,n)<12-46>(K,d)<27>(P,a)KV<1>(I,l)D(D p)(E,v)T<3>W<1>(L,v)L(M,l)(E,d)<02>D(K,e)<1>(K,t)<1>(E,a)(W,k)(W,l)<1>(E,k)ER<2>F<1>(G,t)R<1>(D,n)(S,l)FI(A,n)RM(H,r)<1>(V,l)QG(D,n)R<1>F<1>(P,q)WKGSVVDSV(I,v)GVFLTQN(V,t)D(H,y)(L,s)SS(S,n)A(F,y)M<1>(L,v)A(A,s)<1>FP<0-16>(P,v)<6-15>(S,h)<3>(E,d)<10-</pre>

- 24 > (S,t) < 1 > (S,e) < 6 > (K,n) < 8 55 > (E,i) < 8 9 > (I,v) < 1 > (N,s) < 1 4 > (E,d) < 1 > (E,s) < 4 > (Q,1) < 0 11 > (D,h) < 1 > (F,m) < 5 > (Q,n) < 0 3 > (G,e) < 2 > (G,d) S < 1 > (K,d) < 7 11 > (T,m) < 2 > (V,1) < 3 > (S,q) < 6 10 > (S,e) < 2 3 > (S,v) < 19 25 > (T,s) < 16 28 > (R,s) < 2 6 > (T,p) < 5 > (P,k) < 10 > (Q,e) < 4 > (D,s) < 1 4 > (S,r) < 5 > (D,p) < 3 > (N,d) < 3 > (P,y) < 2 > (F,s) < 1 > (R,k) < 1 > (G,s) < 1 > (S,a) (V,r) (P,e) < 3 > (T,s) < 1 > (R,k) < 1 >
- 35 L,i)GLKS(V,a)ECVRLL<1>L(H,k)<2>AFPVDTNVGRI(A,c)VR(M,1)G(W,1)VPL(Q,e)PLP<2>(L,v)Q(L,m)H(L,q)L(E,f)<1>YP<1>(L,m)(E,d)(S,n)(I,v)QK(F,y)LWPRLCKL(D,p)Q<1>TLYELHY(Q,h)(L,m)ITFGK<0-2>FCTK<2>PNCNACPM(R,k)<0-2>EC(R,k)(H,y)(F,y)(A,s)SA<1>(A,v)<0-10>S(A,s)(R,k)<1>(A,1)L(P,e)<1>(P,t)(E,q)<7-16>P(I,1)(I,v)E(E,f)P<1>(S,t)P<2-5>E<0-15>(D,a)IE(D,e)<4-23>(I,v)P<1>I<1>(L,f)(N,d)<8-
- 40 17>(S,a)<1>(A,d)LV<8>(I,1)P<25>(K,r)(L,m)K<4>LRTEH</br>
 >V(Y,f)(E,v)LPD<1>H<1>(L,i)L(E,k)<1>(D,e)D(P,i)<2>YLL(A,s)
)IW(T,q)P(G,d)(E,g)<6-3>(P,s)<3>C<610>(M,1)C<4>C<2>C<3>(R,k)E<5>(V,f)RGT(L,i)L<022>(L,v)FADH<1>(S,t)(S,r)<2>PI<3>(R,t)<3>(W,k)<1>L<1>(R,k)R<4>G(T,s)(S,t)<2>(S,t)<2>(S,t)<1>(S,t)<2>(S,t)<2>(S,t)<1>(S,t)<2>(S,t)<1>(S,t)<2>(S,t)<1>(S,t)<1>(S,t)<1>(S,t)<1>(S,t)<1>(S,t)<1>(S,t)<1>(S,t)<1>(S,t)<1>(S,t)<1>(S,t)<1>(S,t)<1>(S,t)<1>(S,t)<1>(S,t)<1>(S,t)<1>(S,t)<1>(S,t)<1>(S,t)<1>(S,t)<1>(S,t)<1>(S,t)<1>(S,t)<1>(S,t)<1>(S,t)<1>(S,t)<1>(S,t)<1>(S,t)<1>(S,t)<1>(S,t)<1>(S,t)<1>(S,t)<1>(S,t)<1>(S,t)<1>(S,t)<1>(S,t)<1>(S,t)<1>(S,t)<1>(S,t)<1>(S,t)<1>(S,t)<1>(S,t)<1>(S,t)<1>(S,t)<1>(S,t)<1>(S,t)<1>(S,t)<1>(S,t)<1>(S,t)<1>(S,t)<1>(S,t)<1>(S,t)<1>(S,t)<1>(S,t)<1>(S,t)<1>(S,t)<1>(S,t)<1>(S,t)<1>(S,t)<1>(S,t)<1>(S,t)<1>(S,t)<1>(S,t)<1 (S,t)<1 (S,t)<

10

15

20

25

30

) I (F, c) (R, k) (G, l) L<1> (T, v) <2>I<2> (C, n) F(W,q) <1>G(F,y) (V,l) C(V,l) R<1>F(E,d) <3> (R,g) <1>P(R,k) <1>L<2> (R,h) LH<2> (A,v) SK

DMT domain A corresponds to amino acid positions 697 through 796 of SEQ ID NO:2. DMT domain B corresponds to amino acid positions 1192 through 1404 of SEQ ID NO:2. DMT domain C corresponds to amino acid positions 1452 through 1722 of SEQ ID NO:2. The consensus sequence provides amino acid sequences by position using single letter amino acid abbreviations. Numbers in carrots ("<" or ">") refer to amino acid positions where there is no consensus and which therefore, can be any amino acid. Amino acid abbreviations in parentheses indicate alternative amino acids at the same position. Capitalized letters refer to predominant consensus amino acids and lower case letters refer to amino acids that are commonly found in DMT sequences, but are not predominant. Thus, it is a simple matter to identify whether any particular nucleic acid sequence is a DMT nucleic acid and/or encodes a DMT polypeptide.

The structure of full-length DMT polypeptides comprises the following domains and regions. These regions are generally described with reference to SEQ ID NO:2. First, as described above, domain B DMT polypeptides can comprise a bipartite nuclear localization signal (e.g., amino acid positions 43-60 and 61-78 in SEQ ID NO:2) comprised of basic amino acids. Amino acids 36-91 are homologous to human G/T mismatch-specific thymine DNA glycosylase (Genbank accession number AAC50540.1), which has 5-methylcytosine glycosylase activity (Zhu *et al.*, *Nuc. Acids Res.* 28:4157-4165 (2000)). DMT polypeptides also contain a leucine zipper sequence (e.g., positions 1330-1351 of SEQ ID NO:2), that can be involved in protein-protein interactions as well as DNA binding. In addition, the amino portion of the DMT polypeptide (amino acids 43-78) is generally basic, similar to histone H1. Thus, without intending to limit the scope of the invention, it is believed this basic portion of DMT facilitates interactions with DNA and/or chromatic proteins.

In addition, amino acids 1-800 is related to the beta subunit of bacterial DNA-dependent RNA polymerases. Without intending to limit the scope of the invention, it is believed the RNA polymerase-like domain facilitates interaction of DMT with DNA.

Amino acids 1167-1368 is related to proteins in the HhH-GPD superfamily. Amino acids 1,271 to 1,304 correspond to the conserved HhH-GPD motif. The corresponding DMT sequence is

DKAKDYLLSIRGLGLKSVECVRLLTLHNLAFPVD. Secondary structure prediction

10

15

20

25

30

(Jpred program) indicates that DMT has two alpha-helices (1,271 - 1,279 and 1,286 to 1,295) that correspond to the conserved alphaK and alphaL helices in the HhH-GPD motif of the crystallized hOGG1 DNA repair protein (Bruner et al *Nature* 403:859-866 (2000)). In between the two helices (1280 to 1285), is a hairpin with conserved glycines (G1282 and G1284). Amino acids 1286 to 1295 are related to the alphaL helix of hOGG1, which contacts the DNA backbone (Bruner *et al Nature* 403:859-866 (2000). Thus, without intending to limit the scope of the invention, it is believed this region of DMT contacts the DNA. The catalytic lysine (K1286) and aspartic acid (D1304) residues are conserved in the HhH-GPD motif of DMT. Without intending to limit the scope of the invention, by analogy to hOGG1, K1286 is predicted to displace the modified base and to promote conjugate elimination of the 3'-phosphodiester bond. Without intending to limit the scope of the invention, by analogy to hOGG1, D1304 is believed to assist the reaction by transferring protons to and from K1286.

DMT nucleic acids are a new class of plant regulatory genes that encode polypeptides with sequence identity to members of the endonuclease III genes found in a diverse collection of organisms. Endonuclease III is implicated in various DNA repair reactions. Thus proteins related to endonuclease III are likely to have a chromosomal function. DMT (SEQ ID NO:1) is most related to endonuclease III from Deinococcus radiodurans Genbank Accession No. AE002073 (see, e.g., White, O. et al. Science 286:1571-1577 (1999)). DMT polypeptides have glycosylase activity (i.e., the capability to cleave the base portion of a nucleotide from the sugar portion). More particularly, DMT polypeptides have demethylase activity, and in more preferred embodiments, have 5-methylcytosine glycosylase activity. Demethylation activity can be assayed in vivo by expressing a candidate polypeptide in the nucleus of a cell and then assaying for a change in methylation of the cell's DNA. See, e.g., Vong, et al., Science 260:1926-1928 (1993). Changes in chromosomal methylation can be measured by comparing the ability of methylation sensitive and insensitive endonucleases to cleave DNA from a cell expressing a polypeptide suspected of having demethylase or methylase activity. Alternatively, bisulfate sequencing can be used to identify which base pairs are methylated in a DNA sequence. For a discussion of both methods, see Soppe et al., Molec. Cell. 6:791-802 (2000). In vitro assays to measure demethylase activity using labeled substrates are also known to those of skill in the art. See, e.g., Vhu et al., Proc. Natl. Acad. Sci. USA 97:5135-5139 (2000).

10

15

20

25

30

In the case of both expression of transgenes and inhibition of endogenous genes (e.g., by antisense, or sense suppression) one of skill will recognize that the inserted polynucleotide sequence need not be identical, but may be only "substantially identical" to a sequence of the gene from which it was derived. As explained below, these substantially identical variants are specifically covered by the term *DMT* nucleic acid.

In the case where the inserted polynucleotide sequence is transcribed and translated to produce a functional polypeptide, one of skill will recognize that because of codon degeneracy a number of polynucleotide sequences will encode the same polypeptide. These variants are specifically covered by the terms "DMT nucleic acid". In addition, the term specifically includes those sequences substantially identical (determined as described below) with a DMT polynucleotide sequence disclosed here and that encode polypeptides that are either mutants of wild type DMT polypeptides or retain the function of the DMT polypeptide (e.g., resulting from conservative substitutions of amino acids in the DMT polypeptide). In addition, variants can be those that encode dominant negative mutants as described below.

Two nucleic acid sequences or polypeptides are said to be "identical" if the sequence of nucleotides or amino acid residues, respectively, in the two sequences is the same when aligned for maximum correspondence as described below. The terms "identical" or percent "identity," in the context of two or more nucleic acids or polypeptide sequences, refer to two or more sequences or subsequences that are the same or have a specified percentage of amino acid residues or nucleotides that are the same, when compared and aligned for maximum correspondence over a comparison window, as measured using one of the following sequence comparison algorithms or by manual alignment and visual inspection. When percentage of sequence identity is used in reference to proteins or peptides, it is recognized that residue positions that are not identical often differ by conservative amino acid substitutions, where amino acids residues are substituted for other amino acid residues with similar chemical properties (e.g., charge or hydrophobicity) and therefore do not change the functional properties of the molecule. Where sequences differ in conservative substitutions, the percent sequence identity may be adjusted upwards to correct for the conservative nature of the substitution. Means for making this adjustment are well known to those of skill in the art. Typically this involves scoring a conservative substitution as a partial rather than a full mismatch, thereby increasing the percentage sequence identity. Thus, for example, where an identical amino acid is given a score of 1 and a non-conservative substitution is given a

10

15

20

25

30

score of zero, a conservative substitution is given a score between zero and 1. The scoring of conservative substitutions is calculated according to, e.g., the algorithm of Meyers & Miller, *Computer Applic. Biol. Sci.* 4:11-17 (1988) e.g., as implemented in the program PC/GENE (Intelligenetics, Mountain View, California, USA).

The phrase "substantially identical," in the context of two nucleic acids or polypeptides, refers to a sequence or subsequence that has at least 40% sequence identity with a reference sequence. Alternatively, percent identity can be any integer from 40% to 100%. More preferred embodiments include at least: 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 99% compared to a reference sequence using the programs described herein; preferably BLAST using standard parameters, as described below. This definition also refers to the complement of a test sequence, when the test sequence has substantial identity to a reference sequence.

For sequence comparison, typically one sequence acts as a reference sequence, to which test sequences are compared. When using a sequence comparison algorithm, test and reference sequences are entered into a computer, subsequence coordinates are designated, if necessary, and sequence algorithm program parameters are designated. Default program parameters can be used, or alternative parameters can be designated. The sequence comparison algorithm then calculates the percent sequence identities for the test sequences relative to the reference sequence, based on the program parameters.

A "comparison window", as used herein, includes reference to a segment of any one of the number of contiguous positions selected from the group consisting of from 20 to 600, usually about 50 to about 200, more usually about 100 to about 150 in which a sequence may be compared to a reference sequence of the same number of contiguous positions after the two sequences are optimally aligned. Methods of alignment of sequences for comparison are well-known in the art. Optimal alignment of sequences for comparison can be conducted, e.g., by the local homology algorithm of Smith & Waterman, *Adv. Appl. Math.* 2:482 (1981), by the homology alignment algorithm of Needleman & Wunsch, *J. Mol. Biol.* 48:443 (1970), by the search for similarity method of Pearson & Lipman, *Proc. Nat'l. Acad. Sci. USA* 85:2444 (1988), by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Dr., Madison, WI), or by manual alignment and visual inspection.

10

15

20

One example of a useful algorithm is PILEUP. PILEUP creates a multiple sequence alignment from a group of related sequences using progressive, pairwise alignments to show relationship and percent sequence identity. It also plots a tree or dendogram showing the clustering relationships used to create the alignment. PILEUP uses a simplification of the progressive alignment method of Feng & Doolittle, J. Mol. Evol. 35:351-360 (1987). The method used is similar to the method described by Higgins & Sharp, CABIOS 5:151-153 (1989). The program can align up to 300 sequences, each of a maximum length of 5,000 nucleotides or amino acids. The multiple alignment procedure begins with the pairwise alignment of the two most similar sequences, producing a cluster of two aligned sequences. This cluster is then aligned to the next most related sequence or cluster of aligned sequences. Two clusters of sequences are aligned by a simple extension of the pairwise alignment of two individual sequences. The final alignment is achieved by a series of progressive, pairwise alignments. The program is run by designating specific sequences and their amino acid or nucleotide coordinates for regions of sequence comparison and by designating the program parameters. For example, a reference sequence can be compared to other test sequences to determine the percent sequence identity relationship using the following parameters: default gap weight (3.00), default gap length weight (0.10), and weighted end gaps.

)

30

Another example of algorithm that is suitable for determining percent sequence identity and sequence similarity is the BLAST algorithm, which is described in Altschul et al., J. Mol. Biol. 215:403-410 (1990). Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information (http://www.ncbi.nlm.nih.gov/). This algorithm involves first identifying high scoring sequence pairs (HSPs) by identifying short words of length W in the query sequence, which either match or satisfy some positive-valued threshold score T when aligned with a word of the same length in a database sequence. T is referred to as the neighborhood word score threshold (Altschul et al, supra). These initial neighborhood word hits act as seeds for initiating searches to find longer HSPs containing them. The word hits are extended in both directions along each sequence for as far as the cumulative alignment score can be increased. Extension of the word hits in each direction are halted when: the cumulative alignment score falls off by the quantity X from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters W, T, and X determine the sensitivity and speed of the

SVB B)

5

10

15

20

25

30

alignment. The BLAST program uses as defaults a wordlength (W) of 11, the BLOSUM62 scoring matrix (See Henikoff & Henikoff, Proc. Natl. Acad. Sci. USA 89:10915 (1989)) alignments (B) of 50, expectation (E) of 10, M=5, N=-4, and a comparison of both strands.

The BLAST algorithm also performs a statistical analysis of the similarity between two sequences (see, e.g., Karlin & Altschul, Proc. Nat'l. Acad. Sci. USA 90:5873-5787 (1993)). One measure of similarity provided by the BLAST algorithm is the smallest sum probability (P(N)), which provides an indication of the probability by which a match between two nucleotide or amino acid sequences would occur by chance. For example, a nucleic acid is considered similar to a reference sequence if the smallest sum probability in a comparison of the test nucleic acid to the reference nucleic acid is less than about 0.2, more preferably less than about 0.01, and most preferably less than about 0.001.

"Conservatively modified variants" applies to both amino acid and nucleic acid sequences. With respect to particular nucleic acid sequences, conservatively modified variants refers to those nucleic acids which encode identical or essentially identical amino acid sequences, or where the nucleic acid does not encode an amino acid sequence, to essentially identical sequences. Because of the degeneracy of the genetic code, a large number of functionally identical nucleic acids encode any given protein. For instance, the codons GCA, GCC, GCG and GCU all encode the amino acid alanine. Thus, at every position where an alanine is specified by a codon, the codon can be altered to any of the corresponding codons described without altering the encoded polypeptide. Such nucleic acid variations are "silent variations," which are one species of conservatively modified variations. Every nucleic acid sequence herein which encodes a polypeptide also describes every possible silent variation of the nucleic acid. One of skill will recognize that each codon in a nucleic acid (except AUG, which is ordinarily the only codon for methionine) can be modified to yield a functionally identical molecule. Accordingly, each silent variation of a nucleic acid which encodes a polypeptide is implicit in each described sequence.

As to amino acid sequences, one of skill will recognize that individual substitutions, deletions or additions to a nucleic acid, peptide, polypeptide, or protein sequence which alters, adds or deletes a single amino acid or a small percentage of amino acids in the encoded sequence is a "conservatively modified variant" where the alteration results in the substitution of an amino acid with a chemically similar amino acid.

20

25

30

Conservative substitution tables providing functionally similar amino acids are well known in the art.

The following six groups each contain amino acids that are conservative substitutions for one another:

- 5 1) Alanine (A), Serine (S), Threonine (T);
 - 2) Aspartic acid (D), Glutamic acid (E);
 - 3) Asparagine (N), Glutamine (Q);
 - 4) Arginine (R), Lysine (K);
 - 5) Isoleucine (I), Leucine (L), Methionine (M), Valine (V); and
- 10 6) Phenylalanine (F), Tyrosine (Y), Tryptophan (W). (see, e.g., Creighton, Proteins (1984)).

An indication that two nucleic acid sequences or polypeptides are substantially identical is that the polypeptide encoded by the first nucleic acid is immunologically cross reactive with the antibodies raised against the polypeptide encoded by the second nucleic acid. Thus, a polypeptide is typically substantially identical to a second polypeptide, for example, where the two peptides differ only by conservative substitutions. Another indication that two nucleic acid sequences are substantially identical is that the two molecules or their complements hybridize to each other under stringent conditions, as described below.

The phrase "selectively (or specifically) hybridizes to" refers to the binding, duplexing, or hybridizing of a molecule only to a particular nucleotide sequence under stringent hybridization conditions when that sequence is present in a complex mixture (e.g., total cellular or library DNA or RNA).

The phrase "stringent hybridization conditions" refers to conditions under which a probe will hybridize to its target subsequence, typically in a complex mixture of nucleic acid, but to no other sequences. Stringent conditions are sequence-dependent and will be different in different circumstances. Longer sequences hybridize specifically at higher temperatures. An extensive guide to the hybridization of nucleic acids is found in Tijssen, *Techniques in Biochemistry and Molecular Biology--Hybridization with Nucleic Probes*, "Overview of principles of hybridization and the strategy of nucleic acid assays" (1993). Generally, highly stringent conditions are selected to be about 5-10°C lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength pH. Low stringency conditions are generally selected to be about 15-30°C below the T_m. The

10

15

20

25

30

T_m is the temperature (under defined ionic strength, pH, and nucleic concentration) at which 50% of the probes complementary to the target hybridize to the target sequence at equilibrium (as the target sequences are present in excess, at T_m, 50% of the probes are occupied at equilibrium). Stringent conditions will be those in which the salt concentration is less than about 1.0 M sodium ion, typically about 0.01 to 1.0 M sodium ion concentration (or other salts) at pH 7.0 to 8.3 and the temperature is at least about 30°C for short probes (e.g., 10 to 50 nucleotides) and at least about 55°C, sometimes 60°C, and sometimes 65°C for long probes (e.g., greater than 50 nucleotides). Stringent conditions may also be achieved with the addition of destabilizing agents such as formamide. For selective or specific hybridization, a positive signal is at least two times background, preferably 10 time background hybridization.

Nucleic acids that do not hybridize to each other under stringent conditions are still substantially identical if the polypeptides which they encode are substantially identical. This occurs, for example, when a copy of a nucleic acid is created using the maximum codon degeneracy permitted by the genetic code. In such cased, the nucleic acids typically hybridize under moderately stringent hybridization conditions.

In the present invention, genomic DNA or cDNA comprising *DMT* nucleic acids of the invention can be identified in standard Southern blots under stringent conditions using the nucleic acid sequences disclosed here. For the purposes of this disclosure, suitable stringent conditions for such hybridizations are those which include a hybridization in a buffer of 40% formamide, 1 M NaCl, 1% SDS at 37°C, and at least one wash in 0.2X SSC at a temperature of at least about 50°C, usually about 55°C to about 60°C and sometimes 65°C, for 20 minutes, or equivalent conditions. A positive hybridization is at least twice background. Those of ordinary skill will readily recognize that alternative hybridization and wash conditions can be utilized to provide conditions of similar stringency.

A further indication that two polynucleotides are substantially identical is if the reference sequence, amplified by a pair of oligonucleotide primers, can then be used as a probe under stringent hybridization conditions to isolate the test sequence from a cDNA or genomic library, or to identify the test sequence in, e.g., a northern or Southern blot.

10

15

20

25

30

DETAILED DESCRIPTION

This invention provides molecular strategies for controlling plant development, including methylation of chromosomal DNA, endosperm development and flowering time.

Reproduction in flowering plants involves two fertilization events in the haploid female gametophyte. One sperm nucleus fertilizes the egg to form the embryo. A second sperm nucleus fertilizes the central cell to form the endosperm, a unique tissue that supports the growth of the embryo. Fertilization also activates maternal tissue differentiation, the ovule integuments form the seed coat and the ovary forms the fruit.

The present invention is based, at least in part, on the discovery of a set of female-gametophytic mutations and the subsequent cloning of the gene involved, termed *DEMETER* (*DMT*), formally known as *ATROPOS* (*ATR*). Two mutant alleles of *DMT* disclosed here were created using a T-DNA tag, thereby disrupting an exon of the gene. The *dmt* mutations affect endosperm production, allowing for increased endosperm development. Generally, the mutant *dmt* alleles are not transmitted by the female gametophyte. Inheritance of a mutant *dmt* allele by the female gametophyte usually results in embryo abortion and endosperm overproduction, even when the pollen bears the wild-type *DMT* allele.

In contrast, transmission of *dmt* mutant alleles through the male gametophyte (*i.e.*, pollen) is ecotype-dependent in *Arabidopsis*. For instance, in some ecotypes (e.g., Columbia), transmission of *dmt* mutant alleles is less than 50%. However, in Landsberg *erecta*, transmission is almost normal.

DMT is a repressor of endosperm both before and after fertilization. DMT is both necessary and sufficient for MEDEA transcription. DMT is related to 5-methylcytosine glycosylases. DMT regulates transcription of specific target genes (i.e., MEA) by a demethylation mechanism. DMT is also required for maintaining the proper global pattern of methylation of chromosomal DNA in cells.

The isolated sequences prepared as described herein, can be used in a number of techniques, for example, to suppress or enhance endogenous DMT gene expression. Modulation of *DMT* gene expression or DMT activity in plants is particularly useful, for example, in producing embryo-less or embryo-reduced seed, seed with increased endosperm, as part of a system to generate seed, to modulate time to flowering, organ identity, size and/or number,meristem size or activity in plants, or to modulate methylation, and thus gene expression in plants. Another use is the expression of DMT

polynucleotides in animal cells, for instance as a DNA repair enzyme useful in preventing the unnatural proliferation of cells (including cancer) due to chromosomal lesions. *See*, *e.g.*, Bruner, *et al.*, *Nature* 403:859 (2000).

As described in more detail below, reduction of expression of DMT in plants results in a number of diverse phenotypes. Without intending to limit the invention to particular embodiments, it is belived that some of the phenotypes that are generated in plants are epigenetic mutations, i.e., effects due to differences in the methylation state of the chromosome that result in altered gene expression. Thus, DMT provides a powerful tool to develop any number of plant lines with a variety of desired phenotypes.

10

15

5

Isolation of DMT nucleic acids

Generally, the nomenclature and the laboratory procedures in recombinant DNA technology described below are those well known and commonly employed in the art. Standard techniques are used for cloning, DNA and RNA isolation, amplification and purification. Generally enzymatic reactions involving DNA ligase, DNA polymerase, restriction endonucleases and the like are performed according to the manufacturer's specifications. These techniques and various other techniques are generally performed according to Sambrook *et al.*, *Molecular Cloning - A Laboratory Manual*, Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, (1989).

20

25

The isolation of *DMT* nucleic acids may be accomplished by a number of techniques. For instance, oligonucleotide probes based on the sequences disclosed here can be used to identify the desired gene in a cDNA or genomic DNA library. To construct genomic libraries, large segments of genomic DNA are generated by random fragmentation, e.g. using restriction endonucleases, and are ligated with vector DNA to form concatemers that can be packaged into the appropriate vector. To prepare a cDNA library, mRNA is isolated from the desired organ, such as ovules, and a cDNA library which contains the *DMT* gene transcript is prepared from the mRNA. Alternatively, cDNA may be prepared from mRNA extracted from other tissues in which *DMT* genes or homologs are expressed.

30.

The cDNA or genomic library can then be screened using a probe based upon the sequence of a cloned *DMT* gene disclosed here. Probes may be used to hybridize with genomic DNA or cDNA sequences to isolate homologous genes in the same or different plant species. Alternatively, antibodies raised against a DMT polypeptide can be used to screen an mRNA expression library.

Alternatively, the nucleic acids of interest can be amplified from nucleic acid samples using amplification techniques. For instance, polymerase chain reaction (PCR) technology can be used to amplify the sequences of the DMT genes directly from genomic DNA, from cDNA, from genomic libraries or cDNA libraries. PCR and other in vitro amplification methods may also be useful, for example, to clone nucleic acid sequences that code for proteins to be expressed, to make nucleic acids to use as probes for detecting the presence of the desired mRNA in samples, for nucleic acid sequencing, or for other purposes. For a general overview of PCR see PCR Protocols: A Guide to Methods and Applications. (Innis, M, Gelfand, D., Sninsky, J. and White, T., eds.),

Academic Press, San Diego (1990).

Appropriate primers and probes for identifying DMT sequences from plant tissues are generated from comparisons of the sequences provided here with other related genes. For instance, DMT can be compared to the other endonuclease III genes, such as Genbank Accession No. AE002073. Using these techniques, one of skill can identify conserved regions in the nucleic acids disclosed here to prepare the appropriate primer and probe sequences. Primers that specifically hybridize to conserved regions in DMT genes can be used to amplify sequences from widely divergent plant species. Appropriate primers for amplification of the genomic region or cDNA of DMT include the following primers:

20

25

30

5

10

15

Xba-SKEN-7; CCTCTAGAGGAATTGTCGGCAAAATCGAG

SKB-8; GGAGAGACGGTTATTGTCAACC

SKB-7; AAAAGTCTACAAGGGAGAGAGAGT

SKB-5; GTAGATGTACATACGTACC

SKEN-8; CCATCCTCCAACAAGTAACAATCCACTC

SKB-6; CACTGAGATTAATTCTTCAGACTCG

SKEN-3.5; CTCAGGCGAGTCAATGCCGGAGAACAC

SKEN-3; CGAGGGCTGATCCGGGGGATAGATATTTT

SKEN-2; CCCCGGATCAGCCCTCGAATTC

SKEN-1; CCCCTGTCTACAAATTCACCACCTGG

SKEL-4; CTGACCCAACTGCTTCTCTTC

skes1.5; TCACCTGTTCTGAACAGACTGG

SKES-1.4; CAGCAGACGAGTCCATAATGCTCTGC

SKES-2.4; GTTTGCCTTCCACGACCACC

SKES-1; G\(\psi\)AAGCCACGCAAAGCTGCAACTCAGG

35 SKES-2.45; GAGTTGCAGCTTTGCGTGGCTTCC

	SKES2.5; TTCAGACTCAGAGTCACCTTGC
	SKES-2; ACCAGCAGCCTTGCTTGGCC
	SKES-3; CATGCCAGAGAAGCAGGGCTCC
	SKES3.5; CGATGATACTGTCTCTCGAGC
5	SKES-6; CCTCCGCCTGCTCATGCCTCAG
	SKEN-4; GTCCATCAGGAGAACTTCTGTGTCAGGAT
	SKES-4; GGGAACAAGTGCACCATCTCC
	SKEN-6; GCTCTCATAGGGAACAAGTGCACCATCTC
	skes-5; cgctcgcatgcacctggtac
10	SKB-1; GGAGGGAATCGAGCAGCTAGAG
	SKB-2; GAGCAGCTAAGGGACTGTTCAAACTC
	SKB-3; CCAGGAATGGGATTGTCCGG
	3' RACE-2; CTTGGACGGCGCTTGAGGAACC
	3' RACE-1; GCCTACAAGCCAGTGGGATAG
15	cDNA-1; GCCAAGGACTATCTCTTGAGC
	SKB-4; GGATGGACTCGAGCACTGGG
	SKE2.2-4; AGAGGAGAGTGCAGACACTTTG
	cDNA-3; GAGGACCCTGACGAGATCCCAAC
	cDNA-9; CCATGTGTTCCCGTAGAGTCATTCC
20	2.2+SKE-1; ATGGAGCTCCAAGAAGGTGACATG
	cDNA-5; CAGAAGTGTGGAGGGAAAGCGTCTGGC
	cDNA-4; CCCTCAGACTGTTACACTCAGAAC
	cDNA-2; CCCCTTGAGCGGAAAACTTCCTCTCATGGC
	cDNA-7; GGAAAGGATTCGTATGTGTCCGTGG
25	SKEN-5; GCAATGCGTTTGCTTTCCAGTCATCT
	cDNA-6; GAGGAGAGCAGAGAAGCAATGCGTTTGC
	cDNA-8; GTTAGAGAGAAAATAAATAACCC
	2.2+SKE-3; CCGTAAACAACACCGGATACAC

35

The amplification conditions are typically as follows. Reaction components: 10 mM Tris-HCl, pH 8.3, 50 mM potassium chloride, 1.5 mM magnesium chloride, 0.001% gelatin, 200 μ M dATP, 200 μ M dCTP, 200 μ M dGTP, 200 μ M dTTP, 0.4 μ M primers, and 100 units per ml Taq polymerase. Program: 96 C for 3 min., 30 cycles of 96 C for 45 sec., 50 C for 60 sec., 72 for 60 sec, followed by 72 C for 5 min.

Standard nucleic acid hybridization techniques using the conditions disclosed above can then be used to identify full-length cDNA or genomic clones.

10

15

25

30

Alternatively, a number of methods for designing modifications of polynucleotide sequences are known to those of skill in the art. For example, oligonucleotide directed mutagenesis can be used to introduce site-specific mutations in a nucleic acid sequence of interest. Examples of such techniques are found in the references above and, e.g., in Reidhaar-Olson et al. Science, 241:53-57 (1988) and Ausubel et al. Similarly, gene shuffling (Stemmer Proc. Natl. Acad. Sci. USA 91:10747-10751(1994); Ostermeier et al. Proc. Natl. Acad. Sci. USA, 96: 3562-67(1999))) can be used to introduce variation into one or more DMT sequences or subsequences. For example, orthologous (between species) or homologous (within a species) DMT nucleic acids can be interchanged, combined or shuffled to produce novel variations within the scope of the invention.

Additionally, error prone PCR can also be used to introduce variation into a nucleic acid sequence. See, Leung et al. (1989) Technique 1:11-15 and Caldwell et al. (1992) PCR Methods Applic. 2:28-33.

Control of DMT activity or gene expression

Since *DMT* genes are involved in controlling seed, in particular endosperm, development, inhibition of endogenous DMT activity or gene expression is useful in a number of contexts. For instance, reduction of DMT activity can be used for production of seed with enhanced endosperm. By reducing and/or eliminating DMT activity, plants with seed containing increased endosperm can be produced.

Alternatively, substantial inhibition of DMT activity can be used for production of fruit with small and/or degraded seed (referred to here as "seedless fruit") after fertilization. In many plants, particularly dicots, the endosperm is not persistent and eventually is degraded. Thus, in plants of the invention in which DMT activity is inhibited, embryo-less seed do not persist and seedless fruit are produced. For production of dicots with enhanced endosperm, the most beneficial effect may be to reduce, but not eliminate DMT activity. On the other hand, in monocots, which have persistent endosperm, it is advantageous to eliminate DMT activity.

Alternatively, plants of the invention can be used to prevent pre-harvest sprouting in seeds, especially those derived from cereals. In these plants, the endosperm persists and is the major component of the mature seed. Premature growth of embryos in stored grain causes release of degradative enzymes which digest starch and other

10

15

20

25

30

components of the endosperm. Plants of the present invention are useful in addressing this problem because the seeds lack an embryo and thus will not germinate.

Moreover, as discussed herein, time to flowering and DNA methylation can also be modulated by modulating DMT activity in a cell. For example, DMT can be used to modulate the amount of methylated DNA in a cell. Indeed, since expression of many genes is dependent on their methylation state, modulation of DMT activity modulates gene expression in a cell. Examples of genes whose expression is modulated by DMT include *MEDEA*.

One of skill will recognize that a number of methods can be used to modulate DMT activity or gene expression. DMT activity can be modulated in the plant cell at the gene, transcriptional, posttranscriptional, translational, or posttranslational, levels. Techniques for modulating DMT activity at each of these levels are generally well known to one of skill and are discussed briefly below.

Methods for introducing genetic mutations into plant genes are well known. For instance, seeds or other plant material can be treated with a mutagenic chemical substance, according to standard techniques. Such chemical substances include, but are not limited to, the following: diethyl sulfate, ethylene imine, ethyl methanesulfonate and N-nitroso-N-ethylurea. Alternatively, ionizing radiation from sources such as, for example, X-rays or gamma rays can be used.

Alternatively, homologous recombination can be used to induce targeted gene disruptions by specifically deleting or altering the *DMT* gene *in vivo* (see, generally, Grewal and Klar, Genetics 146: 1221-1238 (1997) and Xu et al., Genes Dev. 10:2411-2422 (1996)). Homologous recombination has been demonstrated in plants (Puchta et al., Experientia 50:277-284 (1994), Swoboda et al., EMBO J. 13:484-489 (1994); Offringa et al., Proc. Natl. Acad. Sci. USA 90: 7346-7350 (1993); and Kempin et al. Nature 389:802-803 (1997)).

In applying homologous recombination technology to the genes of the invention, mutations in selected portions of a DMT gene sequences (including 5' upstream, 3' downstream, and intragenic regions) such as those disclosed here are made *in vitro* and then introduced into the desired plant using standard techniques. Since the efficiency of homologous recombination is known to be dependent on the vectors used, use of dicistronic gene targeting vectors as described by Mountford *et al. Proc. Natl. Acad. Sci. USA* 91:4303-4307 (1994); and Vaulont *et al. Transgenic Res.* 4:247-255 (1995) are conveniently used to increase the efficiency of selecting for altered *DMT* gene

10

15

20

25

30

expression in transgenic plants. The mutated gene will interact with the target wild-type gene in such a way that homologous recombination and targeted replacement of the wild-type gene will occur in transgenic plant cells, resulting in suppression of DMT activity.

Alternatively, oligonucleotides composed of a contiguous stretch of RNA and DNA residues in a duplex conformation with double hairpin caps on the ends can be used. The RNA/DNA sequence is designed to align with the sequence of the target *DMT* gene and to contain the desired nucleotide change. Introduction of the chimeric oligonucleotide on an extrachromosomal T-DNA plasmid results in efficient and specific DMT gene conversion directed by chimeric molecules in a small number of transformed plant cells. This method is described in Cole-Strauss *et al. Science* 273:1386-1389 (1996) and Yoon *et al. Proc. Natl. Acad. Sci. USA* 93:2071-2076 (1996).

Gene expression can be inactivated using recombinant DNA techniques by transforming plant cells with constructs comprising transposons or T-DNA sequences. *DMT* mutants prepared by these methods are identified according to standard techniques. For instance, mutants can be detected by PCR or by detecting the presence or absence of *DMT* mRNA, *e.g.*, by Northern blots. Mutants can also be selected by assaying for development of endosperm in the absence of fertilization.

The isolated nucleic acid sequences prepared as described herein, can also be used in a number of techniques to control endogenous *DMT* gene expression at various levels. Subsequences from the sequences disclosed here can be used to control, transcription, RNA accumulation, translation, and the like.

A number of methods can be used to inhibit gene expression in plants. For instance, antisense technology can be conveniently used. To accomplish this, a nucleic acid segment from the desired gene is cloned and operably linked to a promoter such that the antisense strand of RNA will be transcribed. The construct is then transformed into plants and the antisense strand of RNA is produced. In plant cells, it has been suggested that antisense suppression can act at all levels of gene regulation including suppression of RNA translation (*see*, Bourque *Plant Sci.* (*Limerick*) 105:125-149 (1995); Pantopoulos In Progress in Nucleic Acid Research and Molecular Biology, Vol. 48. Cohn, W. E. and K. Moldave (Ed.). Academic Press, Inc.: San Diego, California, USA; London, England, UK. p. 181-238; Heiser *et al. Plant Sci.* (*Shannon*) 127:61-69 (1997)) and by preventing the accumulation of mRNA which encodes the protein of interest, (*see*, Baulcombe *Plant Mol. Bio.* 32:79-88 (1996); Prins and Goldbach *Arch. Virol.* 141:2259-2276 (1996);

10

15

25

30

Metzlaff et al. Cell 88:845-854 (1997), Sheehy et al., Proc. Nat. Acad. Sci. USA, 85:8805-8809 (1988), and Hiatt et al., U.S. Patent No. 4,801,340).

The nucleic acid segment to be introduced generally will be substantially identical to at least a portion of the endogenous *DMT* gene or genes to be repressed. The sequence, however, need not be perfectly identical to inhibit expression. The vectors of the present invention can be designed such that the inhibitory effect applies to other genes within a family of genes exhibiting homology or substantial homology to the target gene.

For antisense suppression, the introduced sequence also need not be full length relative to either the primary transcription product or fully processed mRNA. Generally, higher homology can be used to compensate for the use of a shorter sequence. Furthermore, the introduced sequence need not have the same intron or exon pattern, and homology of non-coding segments may be equally effective. Normally, a sequence of between about 30 or 40 nucleotides and about full length nucleotides should be used, though a sequence of at least about 100 nucleotides is preferred, a sequence of at least about 200 nucleotides is more preferred, and a sequence of about 500 to about 7000 nucleotides is especially preferred.

A number of gene regions can be targeted to suppress *DMT* gene expression. The targets can include, for instance, the coding regions, introns, sequences from exon/intron junctions, 5' or 3' untranslated regions, and the like. In some embodiments, the constructs can be designed to eliminate the ability of regulatory proteins to bind to *DMT* gene sequences that are required for its cell- and/or tissue-specific expression. Such transcriptional regulatory sequences can be located either 5'-, 3'-, or within the coding region of the gene and can be either promote (positive regulatory element) or repress (negative regulatory element) gene transcription. These sequences can be identified using standard deletion analysis, well known to those of skill in the art. Once the sequences are identified, an antisense construct targeting these sequences is introduced into plants to control gene transcription in particular tissue, for instance, in developing ovules and/or seed. In one embodiment, transgenic plants are selected for DMT activity that is reduced but not eliminated.

Oligonucleotide-based triple-helix formation can be used to disrupt DMT gene expression. Triplex DNA can inhibit DNA transcription and replication, generate site-specific mutations, cleave DNA, and induce homologous recombination (see, e.g., Havre and Glazer J. Virology 67:7324-7331 (1993); Scanlon et al. FASEB J. 9:1288-1296 (1995); Giovannangeli et al. Biochemistry 35:10539-10548 (1996); Chan and Glazer J.

10

15

20

25

30

Mol. Medicine (Berlin) 75:267-282 (1997)). Triple helix DNAs can be used to target the same sequences identified for antisense regulation.

Catalytic RNA molecules or ribozymes can also be used to inhibit expression of *DMT* genes. It is possible to design ribozymes that specifically pair with virtually any target RNA and cleave the phosphodiester backbone at a specific location, thereby functionally inactivating the target RNA. In carrying out this cleavage, the ribozyme is not itself altered, and is thus capable of recycling and cleaving other molecules, making it a true enzyme. The inclusion of ribozyme sequences within antisense RNAs confers RNA-cleaving activity upon them, thereby increasing the activity of the constructs. Thus, ribozymes can be used to target the same sequences identified for antisense regulation.

A number of classes of ribozymes have been identified. One class of ribozymes is derived from a number of small circular RNAs which are capable of self-cleavage and replication in plants. The RNAs replicate either alone (viroid RNAs) or with a helper virus (satellite RNAs). Examples include RNAs from avocado sunblotch viroid and the satellite RNAs from tobacco ringspot virus, lucerne transient streak virus, velvet tobacco mottle virus, solanum nodiflorum mottle virus and subterranean clover mottle virus. The design and use of target RNA-specific ribozymes is described in Zhao and Pick *Nature* 365:448-451 (1993); Eastham and Ahlering *J. Urology* 156:1186-1188 (1996); Sokol and Murray *Transgenic Res.* 5:363-371 (1996); Sun *et al. Mol. Biotechnology* 7:241-251 (1997); and Haseloff *et al. Nature*, 334:585-591 (1988).

Another method of suppression is sense cosuppression. Introduction of nucleic acid configured in the sense orientation has been recently shown to be an effective means by which to block the transcription of target genes. For an example of the use of this method to modulate expression of endogenous genes (*see*, Assaad *et al. Plant Mol. Bio.* 22:1067-1085 (1993); Flavell *Proc. Natl. Acad. Sci. USA* 91:3490-3496 (1994); Stam *et al. Annals Bot.* 79:3-12 (1997); Napoli et al., *The Plant Cell* 2:279-289 (1990); and U.S. Patents Nos. 5,034,323, 5,231,020, and 5,283,184).

The suppressive effect may occur where the introduced sequence contains no coding sequence *per se*, but only intron or untranslated sequences homologous to sequences present in the primary transcript of the endogenous sequence. The introduced sequence generally will be substantially identical to the endogenous sequence intended to be repressed. This minimal identity will typically be greater than about 65%, but a higher identity might exert a more effective repression of expression of the endogenous

10

15

20

25

30

sequences. Substantially greater identity of more than about 80% is preferred, though about 95% to absolute identity would be most preferred. As with antisense regulation, the effect should apply to any other proteins within a similar family of genes exhibiting homology or substantial homology.

For sense suppression, the introduced sequence, needing less than absolute identity, also need not be full length, relative to either the primary transcription product or fully processed mRNA. This may be preferred to avoid concurrent production of some plants that are overexpressers. A higher identity in a shorter than full length sequence compensates for a longer, less identical sequence. Furthermore, the introduced sequence need not have the same intron or exon pattern, and identity of non-coding segments will be equally effective. Normally, a sequence of the size ranges noted above for antisense regulation is used. In addition, the same gene regions noted for antisense regulation can be targeted using cosuppression technologies.

In a preferred embodiment, expression of a nucleic acid of interest can be suppressed by the simultaneous expression of both sense and antisense constructs (Waterhouse *et al.*, *Proc. Natl. Acad. Sci. USA* 95:13959-13964 (1998). *See also* Tabara *et al. Science* 282:430-431 (1998).

Alternatively, *DMT* activity may be modulated by eliminating the proteins that are required for *DMT* cell-specific gene expression. Thus, expression of regulatory proteins and/or the sequences that control *DMT* gene expression can be modulated using the methods described here.

Another method is use of engineered tRNA suppression of *DMT* mRNA translation. This method involves the use of suppressor tRNAs to transactivate target genes containing premature stop codons (*see*, Betzner *et al. Plant J.*11:587-595 (1997); and Choisne *et al. Plant J.*11:597-604 (1997). A plant line containing a constitutively expressed *DMT* gene that contains an amber stop codon is first created. Multiple lines of plants, each containing tRNA suppressor gene constructs under the direction of cell-type specific promoters are also generated. The tRNA gene construct is then crossed into the *DMT* line to activate *DMT* activity in a targeted manner. These tRNA suppressor lines could also be used to target the expression of any type of gene to the same cell or tissue types.

DMT proteins may form homogeneous or heterologous complexes *in vivo*. Thus, production of dominant-negative forms of *DMT* polypeptides that are defective in their abilities to bind to other proteins in the complex is a convenient means to inhibit

10

15

20 -

25

30

endogenous DMT activity. This approach involves transformation of plants with constructs encoding mutant DMT polypeptides that form defective complexes and thereby prevent the complex from forming properly. The mutant polypeptide may vary from the naturally occurring sequence at the primary structure level by amino acid substitutions, additions, deletions, and the like. These modifications can be used in a number of combinations to produce the final modified protein chain. Use of dominant negative mutants to inactivate target genes is described in Mizukami *et al. Plant Cell* 8:831-845 (1996).

Another strategy to affect the ability of a DMT protein to interact with itself or with other proteins involves the use of antibodies specific to DMT. In this method cell-specific expression of DMT-specific Abs is used inactivate functional domains through antibody:antigen recognition (see, Hupp et al. Cell 83:237-245 (1995)).

After plants with reduced DMT activity are identified, a recombinant construct capable of expressing low levels of DMT in embryos can be introduced using the methods discussed below. In this fashion, the level of DMT activity can be regulated to produce preferred plant phenotypes. For example, a relatively weak promoter such as the ubiquitin promoter (see, e.g., Garbarino *et al. Plant Physiol.* 109(4):1371-8 (1995); Christensen *et al Transgenic Res.* 5(3):213-8 (1996); and Holtorf *et al. Plant. Mol. Biol.* 29(4):637-46 (1995)) is useful to produce plants with reduced levels of DMT activity or expression. Such plants are useful for producing, for instance, plants that produce seed with enhanced endosperm.

Use of nucleic acids of the invention to enhance DMT gene expression

Isolated sequences prepared as described herein can also be introduced into a plant cell, thereby modulating expression of a particular *DMT* nucleic acid to enhance or increase endogenous gene expression. For instance, without being bound to any theory, in light of DMT's relation to Exonuclease III and DNA glycosylases, applicants believe that DMT binds DNA or chromatin and acts to modulate transcription by modulating the methylation state of DNA. Enhanced expression can therefore be used to control plant morphology by controlling expression of genes under DMT's control, such as *MEDEA*, in desired tissues or cells. Enhanced expression can also be used, for instance, to increase vegetative growth by preventing the plant from setting seed. Where overexpression of a gene is desired, the desired gene from a different species may be used to decrease potential sense suppression effects.

Moreover, as discussed herein, time to flowering and DNA methylation can also be modulated by modulating DMT activity in a cell. For example, increased expression of DMT in a plant results in delayed time to flowering. Similarly, DMT can be used to modulate the amount of methylated DNA in a cell. Indeed, since expression of many genes is dependent on their methylation state, modulation of DMT activity modulates gene expression in a cell. Examples of genes whose expression is modulated by DMT include *MEDEA*.

One of skill will recognize that the polypeptides encoded by the genes of the invention, like other proteins, have different domains that perform different functions. Thus, the gene sequences need not be full length, so long as the desired functional domain of the protein is expressed.

Modified protein chains can also be readily designed utilizing various recombinant DNA techniques well known to those skilled in the art and described in detail, below. For example, the chains can vary from the naturally occurring sequence at the primary structure level by amino acid substitutions, additions, deletions, and the like. These modifications can be used in a number of combinations to produce the final modified protein chain.

Preparation of recombinant vectors

20

25

30

5

10

15

To use isolated sequences in the above techniques, recombinant DNA vectors suitable for transformation of plant cells are prepared. Techniques for transforming a wide variety of flowering plant species are well known and described in the technical and scientific literature. See, for example, Weising *et al. Ann. Rev. Genet.* 22:421-477 (1988). A DNA sequence coding for the desired polypeptide, for example a cDNA sequence encoding a full length protein, will preferably be combined with transcriptional and translational initiation regulatory sequences which will direct the transcription of the sequence from the gene in the intended tissues of the transformed plant.

For example, for overexpression, a plant promoter fragment may be employed which will direct expression of the gene in all tissues of a regenerated plant. Such promoters are referred to herein as "constitutive" promoters and are active under most environmental conditions and states of development or cell differentiation. Examples of constitutive promoters include the cauliflower mosaic virus (CaMV) 35S transcription initiation region, the 1'- or 2'- promoter derived from T-DNA of

10

15

20

25

30

Agrobacterium tumafaciens, and other transcription initiation regions from various plant genes known to those of skill. Such genes include for example, ACT11 from Arabidopsis (Huang et al. Plant Mol. Biol. 33:125-139 (1996)), Cat3 from Arabidopsis (GenBank No. U43147, Zhong et al., Mol. Gen. Genet. 251:196-203 (1996)), the gene encoding stearoyl-acyl carrier protein desaturase from Brassica napus (Genbank No. X74782, Solocombe et al. Plant Physiol. 104:1167-1176 (1994)), GPc1 from maize (GenBank No. X15596, Martinez et al. J. Mol. Biol 208:551-565 (1989)), and Gpc2 from maize (GenBank No. U45855, Manjunath et al., Plant Mol. Biol. 33:97-112 (1997)).

Alternatively, the plant promoter may direct expression of the *DMT* nucleic acid in a specific tissue or may be otherwise under more precise environmental or developmental control. Examples of environmental conditions that may effect transcription by inducible promoters include anaerobic conditions, elevated temperature, or the presence of light. Such promoters are referred to here as "inducible" or "tissue-specific" promoters. One of skill will recognize that a tissue-specific promoter may drive expression of operably linked sequences in tissues other than the target tissue. Thus, as used herein a tissue-specific promoter is one that drives expression preferentially in the target tissue, but may also lead to some expression in other tissues as well.

Examples of promoters under developmental control include promoters that initiate transcription only (or primarily only) in certain tissues, such as fruit, seeds, or flowers. Promoters that direct expression of nucleic acids in ovules, flowers or seeds are particularly useful in the present invention. As used herein a seed-specific promoter is one which directs expression in seed tissues, such promoters may be, for example, ovulespecific (which includes promoters which direct expression in maternal tissues or the female gametophyte, such as egg cells or the central cell), embryo-specific, endospermspecific, integument-specific, seed coat-specific, or some combination thereof. Examples include a promoter from the ovule-specific BEL1 gene described in Reiser et al. Cell 83:735-742 (1995) (GenBank No. U39944). Other suitable seed specific promoters are derived from the following genes: MAC1 from maize (Sheridan et al. Genetics 142:1009-1020 (1996), Cat3 from maize (GenBank No. L05934, Abler et al. Plant Mol. Biol. 22:10131-1038 (1993), the gene encoding oleosin 18kD from maize (GenBank No. J05212, Lee et al. Plant Mol. Biol. 26:1981-1987 (1994)), vivparous-1 from Arabidopsis (Genbank No. U93215), the gene encoding oleosin from *Arabidopsis* (Genbank No. Z17657), Atmyc1 from Arabidopsis (Urao et al. Plant Mol. Biol. 32:571-576 (1996), the 2s seed storage protein gene family from Arabidopsis (Conceicao et al. Plant 5:493-505

10

15

20

25

30

(1994)) the gene encoding oleosin 20kD from *Brassica napus* (GenBank No. M63985), *napA* from *Brassica napus* (GenBank No. J02798, Josefsson *et al. JBL* 26:12196-1301 (1987), the napin gene family from *Brassica napus* (Sjodahl *et al. Planta* 197:264-271 (1995), the gene encoding the 2S storage protein from *Brassica napus* (Dasgupta *et al. Gene* 133:301-302 (1993)), the genes encoding oleosin A (Genbank No. U09118) and oleosin B (Genbank No. U09119) from soybean and the gene encoding low molecular weight sulphur rich protein from soybean (Choi *et al. Mol Gen, Genet.* 246:266-268 (1995)).

In addition, the promoter sequences from the DMT genes disclosed here can be used to drive expression of the DMT polynucleotides of the invention or heterologous sequences. The sequences of the promoters are identified below.

If proper polypeptide expression is desired, a polyadenylation region at the 3'-end of the coding region should be included. The polyadenylation region can be derived from the natural gene, from a variety of other plant genes, or from T-DNA.

The vector comprising the sequences (e.g., promoters or coding regions) from genes of the invention will typically comprise a marker gene which confers a selectable phenotype on plant cells. For example, the marker may encode biocide resistance, particularly antibiotic resistance, such as resistance to kanamycin, G418, bleomycin, hygromycin, or herbicide resistance, such as resistance to chlorosulfuron or Basta.

Promoter and Enhancer Nucleic Acids of the Invention

The present invention provides polynucleotides useful as promoters and enhancers. The invention also provides methods of targeting heterologous polypeptides to a female gametophyte of a plant, including, e.g., the polar nuclei, the eggs and synergids and central cells. Promoter polynucleotides of the invention include, for example, sequences and subsequences of the DMT 5' flanking DNA (SEQ ID NO:3), the 5' UTR region (SEQ ID NO:6) and the 3' flanking region (SEQ ID NO:4). In some embodiments, the promoter sequences are operably linked to the 5' end of the DMT coding region, which is in turn fused to a polynucleotide of interest, typically encoding a polypeptide. An exemplary promoter sequence includes the last 3424 nucleotides of SEQ ID NO:3 linked to the first 1478 nucleotides of SEQ ID NO:5. In some embodiments, a further 444 nucleotides (e.g., the first 444 nucleotides of the DMT coding region) are incorporated into the promoter. In some embodiments, the promoter

10

15

20

25

30

sequences of the invention specifically direct expression of polynucleotides to the female gametophyte and does not direct expression in tissues following fertilization.

Production of transgenic plants

DNA constructs of the invention may be introduced into the genome of the desired plant host by a variety of conventional techniques. For example, the DNA construct may be introduced directly into the genomic DNA of the plant cell using techniques such as electroporation and microinjection of plant cell protoplasts, or the DNA constructs can be introduced directly to plant tissue using ballistic methods, such as DNA particle bombardment.

Microinjection techniques are known in the art and well described in the scientific and patent literature. The introduction of DNA constructs using polyethylene glycol precipitation is described in Paszkowski et al. *Embo J.* 3:2717-2722 (1984). Electroporation techniques are described in Fromm et al. *Proc. Natl. Acad. Sci. USA* 82:5824 (1985). Ballistic transformation techniques are described in Klein et al. *Nature* 327:70-73 (1987).

Alternatively, the DNA constructs may be combined with suitable T-DNA flanking regions and introduced into a conventional Agrobacterium tumefaciens host vector. The virulence functions of the Agrobacterium tumefaciens host will direct the insertion of the construct and adjacent marker into the plant cell DNA when the cell is infected by the bacteria. Agrobacterium tumefaciens-mediated transformation techniques, including disarming and use of binary vectors, are well described in the scientific literature. See, for example Horsch et al. Science 233:496-498 (1984), and Fraley et al. Proc. Natl. Acad. Sci. USA 80:4803 (1983).

Transformed plant cells which are derived by any of the above transformation techniques can be cultured to regenerate a whole plant which possesses the transformed genotype and thus the desired phenotype such as increased seed mass. Such regeneration techniques rely on manipulation of certain phytohormones in a tissue culture growth medium, typically relying on a biocide and/or herbicide marker which has been introduced together with the desired nucleotide sequences. Plant regeneration from cultured protoplasts is described in Evans et al., *Protoplasts Isolation and Culture*, *Handbook of Plant Cell Culture*, pp. 124-176, MacMillilan Publishing Company, New York, 1983; and Binding, *Regeneration of Plants*, *Plant Protoplasts*, pp. 21-73, CRC Press, Boca Raton, 1985. Regeneration can also be obtained from plant callus, explants,

10

15

20

organs, or parts thereof. Such regeneration techniques are described generally in Klee et al. Ann. Rev. of Plant Phys. 38:467-486 (1987).

The nucleic acids of the invention can be used to confer desired traits on essentially any plant. Thus, the invention has use over a broad range of plants, including species from the genera Anacardium, Arachis, Asparagus, Atropa, Avena, Brassica, Citrus, Citrullus, Capsicum, Carthamus, Cocos, Coffea, Cucumis, Cucurbita, Daucus, Elaeis, Fragaria, Glycine, Gossypium, Helianthus, Heterocallis, Hordeum, Hyoscyamus, Lactuca, Linum, Lolium, Lupinus, Lycopersicon, Malus, Manihot, Majorana, Medicago, Nicotiana, Olea, Oryza, Panieum, Pannesetum, Persea, Phaseolus, Pistachia, Pisum, Pyrus, Prunus, Raphanus, Ricinus, Secale, Senecio, Sinapis, Solanum, Sorghum, Theobromus, Trigonella, Triticum, Vicia, Vitis, Vigna, and Zea.

One of skill will recognize that after the expression cassette is stably incorporated in transgenic plants and confirmed to be operable, it can be introduced into other plants by sexual crossing. Any of a number of standard breeding techniques can be used, depending upon the species to be crossed.

Seed obtained from plants of the present invention can be analyzed according to well known procedures to identify plants with the desired trait. If antisense or other techniques are used to control DMT gene expression, Northern blot analysis can be used to screen for desired plants. In addition, the presence of fertilization independent reproductive development can be detected. Plants can be screened, for instance, for the ability to form embryo-less seed, form seed that abort after fertilization, or set fruit in the absence of fertilization. These procedures will depend, part on the particular plant species being used, but will be carried out according to methods well known to those of skill.

25

30

DMT Mutations, Fragments And Fusions

As discussed above, DMT polynucleotides and polypeptides are not limited to the sequences disclosed herein. Those of skill in the art that conservative amino acid substitutions, as well as amino acid additions or deletions may not result in any change in biological activity. Moreover, sequence variants with at least one modulated biological activity of DMT are also contemplated. For example, at least one DMT activity can be increased or decreased by introduction of single or multiple amino acid changes from the sequences disclosed herein. Those of skill in the art will recognize that conservative amino acid substitutions in important functional domains are typically

10

15

20

25

30

useful in generating more active DMT polypeptides. Conversely, non-conservative substitutions of amino acid residues in functional domains, such as the HhH region of DMT (e.g., amino acids 1271-1304 of SEQ ID NO:2) are likely to disrupt at least one biological activity such as DNA binding. In some embodiments, the fragments of the invention consist of about 100, 200, 300 400, 500, 600, 700, 800, 900, or 1000 amino acids.

Alternatively, fragments of the sequences disclosed herein are contemplated. In some preferred embodiments, the polypeptide fragments have at least one biological activity of DMT. For example, amino acid sequences comprising DMT domain B represent polypeptide fragments with glycosylase or demethylase activity. In some embodiments, a fragment comprising amino acids 1167-1404, 1192-1404, 1192-1368 or 1167-1368 of SEQ ID NO:2 have glycosylase activity.

Mutations, fragments and fusions are also useful as dominant negative mutations. For instance, different regions of the DMT protein are responsible for different biological activities. Thus, mutation or deletion of one functional domain can eliminate one but not all activities. For example, mutation or deletion of the DNA binding domain may result in proteins that interact with proteins necessary for DMT function, effectively titrating out those proteins and preventing an active DMT protein from acting. Similarly, DMT fragments comprising the DNA binding portion of the protein with an inactive enzymatic domain or lacking an enzymatic domain are also useful as dominant negative mutants by competing with active DMT polypeptides for DNA binding sites. As described herein, domains of DMT that can be modulated include: the leucine zipper, nuclear localization sequence, HhH domain, the aspartic acid of the GPD domain, as well a DMT domains A, B or C. Without intending to limit the scope of the invention, based on the data provided herein, DMT has glycosylase and demethylase activity and is a DNA repair enzyme.

Targeting the polypeptides of the invention to chromosomal regions

Without intending to limit the scope of the invention, based on the data provided herein, it is believed that DMT has glycosylase and/or demethylase activity and is a DNA repair enzyme. DNA methylation plays an important role in the repression of gene transcription during animal development including embryogenesis, myogenesis and blood cell development. Methylated DNA is recognized by MeCP2 which inturn represses gene transcription by recruiting the Sin3 repressor complex that contains

10

15

20

25

30

catalytically active histone deacetylase (Jones et al. Nature Genetics 19(2):187-191 (1998)). Histone H3 and H4 deacetylation contributes to the formation of transcriptionally inactive chromatin. Thus, DMT can be used for the purpose of modulating the activity of target genes through chromatin architecture in animal cells as well as plant cells. For example, in some embodiments, DMT is used to catalytically remove 5-MeC from target gene DNA in several ways: e.g., (1) by fusing DMT to a sequence specific DNA binding protein, or (2) by fusing DMT to a subunit of the target repressor complex such as MeCP2 or Sin3. When combined with cell, tissue, or developmentally specific promoters DMT can be used to modulate specific sets of target genes.

In addition, reactive oxygen species, partially reduced species that are produced as intermediates of aerobic respiration, are powerful oxidizing agents that escape the mitochondria and attach vial cellular components. Ionizing radiation and other agents that generate free radicals also produce reactive oxygen species that can attack the genome and cause lesions that are thought to have a key role in in causing cancer and ageing. For example, 7,8-dihydro-8-oxoguanine (oxoG) is a very deleterious adduct generated by oxidation of the guanine base in DNA. The oxoG protein can pair with either cytosine or adenine during DNA replication. Thus, oxoG residues in DNA give rise to G/C to T/A transversion mutations. These transversions are common somatic mutations found in human cancers. HhH-GPD enzymes, such as those described herein, represent a defense against oxoG by catalysing the expulsion of the oxoG. Thus, in some embodiments, enhanced DMT activity is a method to reduce the incidence of mutations in animal cells. Also, DMT can be used to catalytically remove oxoG from a target gene by fusing DMT to a sequence specific DNA binding protein. When combined with a cell, tissue, or developmentally specific promoters DMT can be used to modulate repair of target genes.

As described above, the polypeptides of the invention can be targeted to chromosomal regions of interest by linking the polypeptides of the invention, including fragments with demethylase activity, to a DNA-binding domain that binds a target sequence. For example, it is known that an enzyme that methylates DNA (Dam methylase) can be targeted to specific sites in the genome (B.V. Steensel and S. Henikoff, *Nature Biotechnology* 18:424-428 (2000)). Specifically, the methylase was tethered to the DNA-binding domain of GAL4. When recombinant GAL4-methylase protein was expressed in transgenic Drosophila, targeted methylation occurred in a region of a few

25

30

kilobases surrounding the GAL4 DNA binding sequence. In a analogous fashion, DMT, or a portion of DMT that has biological activity (e.g., a portion containing the HhH-GpD motif amino acids such as 1167 to 1368 of SEQ ID NO:2), can be tethered (e.g., as a translational fusion or chemically linked) to proteins that interact at specific sites in the genome. As a result, specific targeted regions of the genome are hypomethylated by DMT. As discussed above, typically hypomethylation promotes transcription of genes (S. E. Jacobsen, *Current Biology* 9, 617 (1999). The invention provides compositions and methods for methylation of a desired area of the chromosome by targeting DMT to those regions. Thus, these embodiments provide additional ways to activate transcription of a desired gene in a targeted chromosomal region.

The following Examples are offered by way of illustration, not limitation.

EXAMPLE

15 **Example 1:**

5

10

This example shows the characterization of *dmt* mutant plants and the isolation of *DMT*.

Arabidopsis plants were transformed by infiltrating them with Agrobacterium containing the SKI15 T-DNA vector (generously provided by D. Weigel (Salk Institute, La Jolla, CA)). T1 seeds were harvested. The SKI15 vector has the bialaphos resistance (BAR) gene that allowed us to directly select transgenic plants in soil after spraying with the commercially available herbicide, Basta. Siliques from approximately 5,000 Basta resistant plants were opened, and those displaying approximately 50% seed abortion were identified.

Two lines, B13 and B33, were identified for further characterization. Genetic analysis of the mutants revealed that the *dmt* mutants were female sterile. Male fertility, however, depended on the genetic background of the mutant alleles. For instance, in the Columbia background, transmission of the *dmt* mutation is less than 50%. However, in the Landsberg *erecta* background, transmission through the male was almost normal.

Molecular analysis confirmed that the two mutations were allelic. For example, both the B13 and B33 alleles carry the SKI15 T-DNA within a DMT exon, confirming that disruption of the *DMT* gene resulted in the observed B13 and B33 phenotypes.

10

15

20

25

5'- and 3'- RACE were used to delineate the 5'- and 3'-ends of the cDNA, respectively. 5'-RACE was carried out using reagents and protocols provided by 5' RACE System for Rapid Amplification of cDNA Ends, Version 2.0, GIBCO BRL, LIFE TECHNOLOGIES, Grand Island, NY and Marathon cDNA Amplification Kit, Clontech, Palo Alto, CA. Final gene specific 5'-RACE primers were SKES-4 (GGGAACAAGTGCACCATCTCC) and SKES3.5 (CGATGATACTGTCTCTTCGAGC). 3'-RACE was carried out using reagents and protocols provided by Marathon cDNA Amplification Kit, Clontech, Palo Alto. Final gene-specific 3' end was obtained from cDNA library screening.

The nucleotide sequence of the genomic copy of *DMT* was also determined (SEQ ID NO:1). The 5'-end of the DMT RNA is located at position 3,425 of SEQ ID NO:1. The position of the 3'-end of the DMT RNA is at position 12,504 of SEQ ID NO:1. The position of the ATG translation initiation codon is at position 4,903 of SEQ ID NO:1. The position of the TAA translation termination codon is at position 12,321 of SEQ ID NO:1.

A portion of the *DMT* polynucleotide sequence, including the first exon, is encompassed by the bacterial artificial chromosome (BAC) clone T9J15TRB. For example, sequences 3820-4299, 4319-4558, 4546-5025 and 9320-9777 of SEQ ID NO:1 were previously determined using the BAC clone as a template. Moreover, a separate independently sequenced region (Bork, C. *et al Gene* 28:147-153 (1998)) also overlaps the *DMT* sequence at positions 11,087 to 12,785 of SEQ ID NO:1.

The predicted DMT protein has 1,729 amino acids. This sequence was compared to known protein sequences using BLAST and revealed homology to several Endonuclease III proteins. The highest homology was to the Endonuclease III protein from *Deinococcus radiodurans*, Genbank Accession No. AE002073 (see, e.g., White, O. et al. Science 286:1571-1577 (1999)). Other DMT motifs include two consecutive nuclear localization signals at positions 43-60 and 61-78 and a leucine zipper at positions 1330-1351.

30 **Example 2:**

This example provides further evidence that mutant phenotypes are caused by loss-of-function mutations.

A new allele, dmt-3, was obtained. The dmt-3 allele was caused by insertion of the simple pD991 T-DNA vector (M. R. Sussman, et al., Plant Physiol.

10

15

20

25

30

124:1465 (2000)) into the 2nd exon of the *DMT* gene. In contrast, the previous two alleles, *dmt-1* and *dmt-2*, were caused by insertion of the activation T-DNA vector, SKI015 vector. The mutant phenotypes generated by all three *dmt* alleles are the same. Because pD991 does not have activation sequences, it suggests that all three mutant alleles are loss-of-function alleles. Consistent with this conclusion, seed abortion can be rescued with a transgene with 3,373 base pairs of 5'-*DMT* flanking sequences plus 1,478 base pairs of 5-'UTR ligated to a cDNA encoding the full-length DMT polypeptide (i.e., *DMTp::DMT*). Thus, *dmt/DMT* heterozygous plants that are hemizygous for the *DMTp::DMT* transgene displayed 25% seed abortion. Control *dmt/DMT* plants displayed 50% seed abortion.

Example 3:

This example shows that DMT is necessary and sufficient for MEA gene expression.

As discussed above, when fertilization of *dmt/dmt* homozygous mutant flowers was prevented, fertilization-independent endosperm development was observed. This is very similar to when fertilization of mutant *mea* flowers is prevented. Thus, before fertilization, both *DMT* and *MEA*, a polycomb protein (T. Kiyosue *et al.*, *Proc. Natl. Acad. Sci. USA* 96:4186 (1999)), prevent the central cell of the female gametophyte from forming an endosperm. This is consistent with *DMT* being a positive regulator of MEDEA (MEA).

As further evidence of this relationship, *MEA* RNA accumulates in immature floral (IF) buds and open flowers (OF). However, in *dmt/dmt* mutant plants there was no detectable *MEA* RNA. Thus, *DMT* is necessary for *MEA* gene expression.

In addition, we have generated plants with a transgene, CaMV::DMT, designed to overexpress DMT. The full-length DMT cDNA was ligated to the constitutive cauliflower mosaic virus promoter, CaMV (S. G. Rogers, H. J. Klee, R. B. Horsch, R. T. Fraley, $Meth\ Enzymol\ 153:253\ (1987)$). In control wild type plants, the DMT and MEA genes were not significantly expressed in the leaf. However, in 35S::DMT plants, both DMT and MEA RNA level increased significantly. This shows that DMT is sufficient to induce MEA gene expression in the leaf.

Example 4:

5

10

15

20

25

This example shows that DMT is a member of the HhH-GPD superfamily of DNA repair enzymes.

A BLAST search, followed by a conserved domain search, revealed that DMT is highly related to the HhH-GPD superfamily of base excision DNA repair proteins (i.e., score of 70.1, E-value of 8e⁻¹³). This family contains a diverse range of structurally related DNA repair proteins. The superfamily is called the HhH-GPD family after its hallmark helix-hairpin-helix and Gly/Pro rich loop followed by a conserved aspartate (S. D. Bruner, *et al.*, *Nature* 403:859 (2000)). This includes endonuclease III (EC:4.9.99.18), 8-oxoguanine DNA glycosylases (i.e., yeast OGG1), the thymine DNA glycosylase of methyl-CPG binding protein MBD4 (B. Hendrich, *et al.*, *Nature* 401:301 (1999)), and DNA-3-methyladenine glycosylase II (EC:3.2.2.21). The predicted amino acid sequence of DMT contains many of the conserved amino acids of this superfamily.

The hallmark of the superfamily of base-excision DNA repair proteins is a helix-hairpin-helix structural element followed by a Gly/Pro-rich loop and a conserved aspartic acid (i.e., HhH-GPD motif). The DMT polypeptide is 1,729 amino acids in length. Amino acids 1,271 to 1,304 correspond to the conserved HhH-GPD motif. The DMT sequence is DKAKDYLLSIRGLGLKSVECVRLLTLHNLAFPVD. The catalytic lysine (K1286) and aspartic acid (D1304) residues are conserved in the HhH-GPD motif of DMT. Secondary structure prediction (Jpred program) indicates that DMT has two alpha-helices (amino acids 1,271 - 1,279 and 1,286 to 1,295) that correspond to the conserved alphaK and alphaL helices in the HhH-GPD motif of the crystallized hOGG1 DNA regair protein (Bruner *et al Nature* 403:859-866 (2000)).

The Arabidopsis *DMT* coding sequences were also used to identify homologous sequences in both public and proprietary databases using both the BLAST and PSI-BLAST computer algorithms. This analysis revealed amino acid sequences from several plant species, including wheat, maize, rice, soybean and Arabidopsis (SEQ ID NOs: 7-29). Based on these sequences, the following consensus sequences for DMT were determined:

DMT Domain A

35

15

20

DMT Domain B

W(D,n)<1>(L,f)R<5>E<3

6>D(S,t)<1>(D,n) (Y,w)<3 R<10>I<2>RG(M,q) (N,f)<2>L(A,s)<1>RI<212>FL<3>V<2>(H,n)G<1>IDLEWLR<2>(P,d) (P,s) (D,h)<1>(A,v)K<1>(Y,f)LL(S,e) (I,f)<1>G(
L,i)GLKS(V,a)ECVRLL<1>L(H,k)<2>AFPVDTNVGRI (A,c)VR(M,1)G(W,1)VPL(Q,e)PLP<2>(L,v)Q
(L,m)H(L,q)L(E,f)<1>YP<1>(L,m) (E,d) (S,n) (I,v)QK(F,y)LWPRLCKL(D,p)Q<1>TLYELHY(Q,h
) (L,m)ITFGK<0-2>FCTK<2>PNCNACPM(R,k)<0-2>EC(R,k) (H,y) (F,y) (A,s)SA<1>(A,v)<010>S(A,s) (R,k)<1>(A,1)L(P,e)<1>(P,t)

10 DMT Domain C.

P(I,1) (I,V) E(E,f) P<1>(S,t) P<2-5>E<0-15>(D,a) IE(D,e) <4-23>(I,V) P<1>I<1>(L,f) (N,d) <8-17>(S,a) <1>(A,d) LV<8>(I,1) P<2-5>(K,r) (L,m) K<4>LRTEH<1>V(Y,f) (E,V) LPD<1>H<1>(L,i) L(E,k) <1>(D,e) D(P,i) <2>YLL(A,s) IW(T,q) P(G,d) (E,g) <6-8>(P,s) <3>C<6-10>(M,1) C<4>C<2>C<3>(R,k) E<5>(V,f) RGT(L,i) L<0-

22>(L,v)FADH<1>(S,t)(S,r)<2>PI<3>(R,t)<3>(W,k)<1>L<1>(R,k)R<4>G(T,s)(S,t)<2>(S,t)
)I(F,c)(R,k)(G,1)L<1>(T,v)<2>I<2>(C,n)F(W,q)<1>G(F,y)(V,1)C(V,1)R<1>F(E,d)<3>(R,g)<1>P(R,k)<1>L<2>(R,h)LH<2>(A,v)SK

The first consensus sequence listed above corresponds to amino acid positions 586 through 937 of SEQ ID NO:2. The second consensus sequence listed above corresponds to amino acid positions 1117 through 1722 of SEQ ID NO:2. The consensus sequence provides amino acid sequences by position using single letter amino acid abbreviations. Numbers in carrots ("<" or ">") refer to amino acid positions where there is no consensus and which therefore, can be any amino acid. Amino acid abbreviations in parentheses indicate alternative amino acids at the same position. Capitalized letters refer to predominant consensus amino acids and lower case letters refer to amino acids that are commonly found in DMT sequences, but are not predominant.

30

35

Example 5:

This example demonstrates the relationship between DNA repair and demethylation.

For many years, attention was focused on the ability of DNA glycosylases to repair DNA. For example, glycosylases are involved in the repair of G/T mismatched bases by depurinating the thymidine base moiety. Recently it was shown that avian (B.

10

15

20

30

Zhu et al., Proc. Natl. Acad. Sci. USA 97:5135 (2000)) and mammalian (B. Zhu et al., Nucl. Acid Res. 28:4157 (2000)). G/T mismatch DNA glycosylases also have 5-methylcytosine-DNA glycosylase activity. That is, these enzymes are demethylases that remove 5-methylcytosine that is later replaced by cytosine. Without intending to limit the scope of the invention, it is believed that as a member of this superfamily, DMT is a demethylase (i.e., 5-methylcytosine glycosylase).

The methylation (i.e., amount of 5-methylcytosine) state of a gene can have a profound effect on its expression. In general, hypomethylation is associated with elevated gene expression, whereas hypermethylation is associated with decreased gene expression (S. E. Jacobsen, *Current Biology* 9:617 (1999)). Thus, it is likely that *DMT* activates *MEA* gene expression by reducing its level of methylation.

Mutations in the *DDM1* gene in Arabidopsis reduce by 70% the overall genome cytosine methylation (E. J. Finnegan, et al., Proc. Natl. Acad. Sci. USA 93:8449 (1996); M. J. Ronemus, et al., Science 273:654 (1996)). Such plants develop a number of phenotypic abnormalities including floral phenotypes (T. Kakutani, et al., Proc. Natl. Acad. Sci. USA 93:12406 (1996)). Similarly, phenotypic abnormalities have been observed developing in dmt/dmt homozygous plants that affect petal number, floral organ fusion, and floral organ identity. Moreover, independent CaMV::DMT transgenic lines that overexpress DMT frequently are late-flowering. This is particularly interesting because late flowering of ddm1 plants was shown to be due to hypomethylation of the FWA gene (W. J. J. Soppe et al., Mol Cell 6:791 (2000)). Thus, without intending to limit the scope of the invention, it is believed that both ddm1 loss-of-function mutations and overexpression of DMT (i.e., CaMV::DMT) may result in genome hypomethylation.

25 **Example 6:**

This example demonstrates targeting gene expression to the female gametophyte using a DMT promoter sequence.

DMT RNA accumulates in many plant organs such as immature flowers, mature flowers, open flowers, stems and to a lesser extent, leaves. To understand the spatial and temporal regulation of DMT RNA accumulation, the expression of the DMT promoter fused to reporter genes was analyzed. We fused 2,282 base pairs of 5'-DMT sequences, the full-length 5'-UTR (1,478 base pairs), 444 base pairs of DMT coding sequences that contain a nuclear localization signal to two reporter genes, the green

10

fluorescent protein (*GFP*; (Y. Niwa, *et al.*, *Plant J.* 18:455 (1999))) and β-glucuronidase (*GUS*; (R. A. Jefferson, T. A. Kavanagh, M. V. Bevan, *EMBO J* 6:3901 (1987))). Reporter gene expression was observed in the developing female gametophyte, in the polar nuclei before they fuse, in the egg and synergids, and in the central cell. Expression was not detected after fertilization. Thus, this promoter is useful for targeting gene expression to the female gametophyte.

It is understood that the examples and embodiments described herein are for illustrative purposes only and that various modifications or changes in light thereof will be suggested to persons skilled in the art and are to be included within the spirit and purview of this application and scope of the appended claims. All publications, patents, and patent applications cited herein are hereby incorporated by reference in their entirety for all purposes.



SEQ ID NO:1 DMT genomic sequence

DMY genomic sequence (12,785 bp)

ATAGAAACGATAACTC

AAGCTT AAAGCTACCAACATCGAATTTAGTAAAAGACCCATGATTTGAAATTGGAATTGTCGG 5 CAAAATCGAGAAGATAT AGAGCCGAGACGGGAACAGTGAAAACCACAAAGCGCGTAAGAATGAAACAGTGGGAGAAGG AAGAGAGAA TCTTACCGAT CATTCGAGGGAAAGATGGGAATCAGAGAAAAATCTGGAAAAAAAGAAATTAAGAGAAAGA GAGAGAAGAAAGTGAGGAG 10 GAGGTTAAAGCAGCAAAT TGTGGAGAGATAAAGÀGAGAGAGAGACTGAGCGAGTCAAGTTCGTCGTCGTCTTTAAAAGAA AGAATCCTATATTTGCCT TTTTCTTTACTACTTTATTTCAGACTATTTGCTTATTTTGCCTCAAACTTTTTTGATTGTCACTT 15 TTCGATCCTAAAGT GTTTGACAATTTACCTGCCTTTTCTCCAAGAAAAATCAGAACAGACCACAGCAAATTTATGTA TTTTCTATTAAAAAAG AAAGAAAGAATTCATATTACTTATAGAAATTAAAAGCTAAGCAGTTGAAAACGTGAAAGCAGA **ATTTCTAAAAAAAAATAGT** 20 AAACTGCTACAAACTTATTTATGTGTATATAACATATCTATAAAGAAACTCAAATATATGATA AATCATTTTAACAAAAT TTCTATGAAATTATAATAAAAAAGTCACTTTTGACACTTAAAAGGTTGACAATAACCGTCTCT CCAAAAAAAATCAAA ACATTTATAATTTCTAAAACTATGGTGTAA\\TTTGCTGAAATCAAAAAGAAAGAAGGATTTC 25 **TATATCATAAGTTTCAT** TATTGTATCAAACTTTCAAATTTCATGTAATTTGAAAGGAAAAAATTAAGATATAATGTTGTT TTTGTTTCTTATGTTA CATTTTCATGGAATATATTCATAACAAAAAATQTATTTTAATATGATGAGAGATTACCATCC AAAAGGTCGAACTTAT 30 ATAAAACAAGTTAATAACTAAACAATACATGTGATCÀCAATCAATGACAGTTTTGATCTTAAA ATAGAAATGATTGAGCA AACCTCAAAAATGTCTTCTTAGGATCACAAAATCTTTCCT\\TAGCTTATTAAAGCCGGGAGTTC **AACTCTCTCTCCCTTG** TAGACTTTTTGTTTTCAAATCTTTTTCTTTCAAAAAATCAATAATTAGTTAATGGGCATAATATT 35 **TGGTTTTAATTAAGT** CCATAGATTTTTTAGGACCATCTCTAATCACGACAAATATCCTAAÅTTGTAACACATTTAAAAC TTAAAAGTATTGCATT

CTTCCTTTATGTAATTC

	TCAAATAGGTTGGTAGT
	TAAAAAAAAAACAAATAGTAGATGTACATACGTACCTTTAAAAATATATACTCATATCGAAAGT
	TTTAAATTTTGCGAAAT
5	TAAATACATITATCTATCAATTAAAATACATTTAATAATGCATAATTCTGTAATATCTATC
	AATTTCCATATAGAAC
	CAAAACAAAATAAACATATCAAATAGTTTTAACTTAACAAAAACGTTAGGGAAAAGTTGACCT
	AACTAGCTTGATTGACG
	TTGAACTTGTCAATGCGAAAGCGATATTTCCAATATATACTACATGTAGTATTATTTAT
10	AGTTTCTAAAAAGGTG
	TTGAGTGGATTGTTACTTGTTGGAGGATGCTATTTTTTCCTTCTTGCCATAATATTTTACGAGTA
	TGGGATAACTACATA
	CTCATGATTATGAAACGCTCACTTTATTTGAAAAACCTCCTAATACACCAAATATGTCACTAGA
	TTCCAAAACGTAGACC
15	AATTGTATCTAATCTCAAATTCTCAATCAAAGTATTAATTTACCGATGGTAAGAAAAGTTAACC
	GATATAATTATCAAAA
	${\bf GAAAGAATAAGTCAACAGATTCTTAATCTCTTTATTTTGGTATATGAACATTTGTACAAAAATC.}$
	TCAAAAGATATGTAAC
	TGTTTAAAATATAAATTCACTGAGATTAATTCTTCAGACTCGTGTTAGCTATAATAATGTCAAG
20	AGTTCTTCTTGTTTCA
	AGGAAAAACCTTAAAGATATGTATATTTTCTGTAATTATGATGATAATTTGCTATTCATTGT
	CACAAACATTACTTTA
	AAAAATCGTATTTTCATTACTACAATGTTGACTAAGAACAAAAATACATTGATTATTGATATAT
	CGTCAACTGAATTTTC
25	TTCCGAGGGATATAATTCTCAAACATAGCAAGAATCTCATAATAATGTTTCGTGACTACCTTTA
	GACGAAATTTTTTAA .
	GATTCGTAACGTGACTTATGGTCTCTTGCTGTGGGGGTCAATGCGAATAAATCTAAATGTATG
	GGAGTCAAATAAAATAC
	CAAGAAAAATAAAGGAGCACCCAATAAACTATATGGGACCAGAAATCCTTTCATTGGTTT
30	AAAATAGGATTATCCCG
	AAAGATGAAGGACTAAATTGAAACTGATTGGGGGTAGGAAGAGATCCGTCACAATCATTAAT
	GGCTTCCACGCGGAAACT
	TGTCGTTTATACAATTTCATTAACTTTCGGGTCGGGTTTATATTCCAAATGGGTCAAATAATAT
	TAGTTTAATACACTAA
35	CGGAGTAATTAATTGGTGACTACAATTTTATCAGTTTGGTGCAATTAGAAACGAACATAGTCG
	TAAAATACGAGTTCGGT
	GTTATACCTTTATTTACGTTAAAAAAAATACGAGAATTTTGTGTCAAATTTCAAATTAATT
	GAATATGGAAATTA
	TTAGATACTCTAGCGAAAATAGTGATTATGAGCGTTTTACAAAAATACGATTTTAGCATTGAA

 $\tt CTTACTCAACAATTAGCCCAAAAAAACATCCATAATGCATTTAAACTAGGAATTTTAACAAAC$

	GGTCAAATGTTGGCATGAAGAAGCAAGTTTGCAACATTAAATTTCATTTAAAAATCGTGTTGA
	CATACTTTAAAATCTAA
	ATATAGGAAGAAGACCAAAACATTAAATTTAGTAAGATTCTAATGAACATTTATAAGTTATAA
•	CTTATAACCAACAAAG
5	TTGGGTTTAGCGTTGTTGCTTTATCTGAAAACTTGCAAACTAAACCATTTTAATAGGACTAATG
	ACAATTAACAACAAAA
•	TACACTTAAGCAACAACGTCCTCGTGAATATAATTTGGGCCTCAGGCCCATATTGCTAACGCC
	AACTGATATTTCACTTT
	ATTCCTTCTCACCACACCACTCTCTCTCTATCTCTATCTCTAACGGCATAGCTGACTCAGTG
10	TTCTCCGGCATTGAC
	TCGCCTGAGAATCAGAAAGCTTAGATCGGTGAGCTTTTAGCTCCATTTTCTGTTTATTTA
	TTATTTCCTTTTTTC
	TCTCTCCCTTTTTTATCTGGAATTTGTTCTGCTAAATTTTCCAGCTGTTACATTTTCCGATCACG
	AGAAGAATCACTGGG
15	TTTTTATGTTAATCAATACATGTTCCTGTTTTCTGATCATAAATCTCAGCTATTAACACCTGATT
	TTGATTCTGCGTAAT
	· AAAAACCTCTGATTTGCTTTTATCTTCACTTTCCCCATAAACATTGCTTACTTTATTCGCTCTTC
	TTTTACCGTTTCCAG
	CTAAAAAATTCTTCGCTATTCAATGTGTTTCTCGTTTTGTTGATGAGAAAAATATCTGACAAAA
20	AATCATTTATTGCATT
	TTATGGTGCAGATTCTTAGTTAATGTCGCCTTCTCTAACCAAGTCAGATTAAAAAAGGAGTGTTC
	GTCCATGTTGCTTTGT
	TTTGGTGTTTTGGAGAGAGTTTTCGGAGAGTTAGGTGAGTGTTATTTGGGGTGAGGTAGTGATA
	AGGTTTGAAGGGGGAGT
25	GATTCATCAAGTGTTTATGAATTCGAGGGCTGATCCGGGGGATAGATA
	GGAGAATCAAACTCAA
	CAAGAGTTCATGGGTTCTTGGATTCCATTTACACCCAAAAAACCTAGATCAAGTCTGATGGTA
	GATGAGAGTGATAAA
	CCAGGATCTAAATGGGTTTCCAGGTGGTGAATTTGTAGACAGGGGATTCTGCAACACTGGTGT
30	GGATCATAATGGGGTTT
	TTGATCATGGTGCTCATCAGGGCGTTACCAACTTAAGTATGATGATCAATAGCTTAGCGGGAT
	CACATGCACAAGCTTGG
	AGTAATAGTGAGAGAGTTTTTGGGCAGGAGTGAGTGACTTCTCCTTTAGCACCAGTTATC
	AGAAACACCACCGGTAA
35	TGTAGAGCCGGTCAATGGAAATTTTACTTCAGATGTGGGTATGGTAAATGGTCCTTTCACCCA
	GAGTGGCACTTCTCAAG
	CTGGCTATAATGAGTTTGAATTGGATGACTTGTTGAATCCTGATCAGATGCCCTTCTCCTTCAC
	AAGCTTGCTGAGTGGT
	GGGGATAGCTTATTCAAGGTTCGTCAATGTGAGTGATCAAATCTATTTTCAGTTTTTTTT
40	CTTTCTTCCGTTCTT





50

CTCGATTCCCTCCAAAATTAAGCAGCAGCCGAGAAGATGAAAGGAATGTTAGAAGCGTAGTT

GTTGAAGATCCAGAAGGA

GAAGTTTCTGGGGTTGA

ATCTACTAAAAACTTA





AACGGCGGCTAATGCG

	TCTCCTTGATGGTGTAAGTCAATTTTTAACTCTCTATACTCGAGTTGTTTCACTTGAGCAACA
	CTGTTTAAAAGTCCT
	CATTTGATAAAATAACAGATGGATAAAAGAGAACCAGATGATCCAAGTCCTTATCTCTTAGCT
	ATATGGACACCAGGTGA
5	GAATAAAACTGCAATGTTTCATTCATGTGTCTACAGTATCAAAGAAAG
	AAAGCATTTGAAATAGA
	GTCGGTTAAATATGAAAGTTTGAATCTGTAAATGAAAGCCGGAACGTAGCATTGGTGGATGTT
	ATATGTAAATTAGTTTT
	TGAGATTGGTCTAATGTAGTTGTTTGACTGCCAGGTGAAACAGCGAATTCGGCACAACCGCCT
10	GAACAGAAGTGTGGAGG
	GAAAGCGTCTGGCAAAATGTGCTTTGACGAGACTTGTTCTGAGTGTAACAGTCTGAGGGAAGC
	AAACTCACAGACAGTTC
	GAGGAACTCTTCTGGTGAGATTATCTTGATCTTTTGTGTTGCTCATGAAAAGGAGAAGTGAGA
	ATACAAGTTTGCTAATA
15	TCATTTTTCGTCATTCACAGATACCTTGTCGGACTGCCATGAGAGGAAGTTTTCCGCTCAACG
	GGACATATTTCCAAGT
	CAACGAGGTTAGATGAAATAAAACTCAAACAGACAGACGAAACATTATTTCTGTTTAGTGTTC
	GTTCTTTATCCTCCTTG
	CCATTTTTTATCTTGCAGTTATTTGCAGACCACGAGTCCAGTCTCAAACCCATCGATGTTCCTA
20	GAGATTGGATATGGGA
	TCTCCCAAGAAGGACTGTTTACTTCGGAACATCAGTAACATCAATATTCAGAGGTAAAAACAT
	TCGTAATAGAGTTAGTT
	AATCAAATGTCCAAAACACAAGAAAGCTTCACCGTCCAATACACAAGAAAGCTTCACCTTCTC
	TTTGCCAAAAAGATCT
25	TAGAATGTTTTGCTGAATTTGTGCAGGTCTTTCAACGGAGCAGATACAGTTCTGCTTTTGGAAA
	GGTAAACGTTAACTTT
	CGACCCAGAGAAATCCGGAAAATCTATTGCTTTGTTCTGATCAATACGTTAAACATATACACA
	CACACTTTACACTTAGG
	ACCAATACTGTTCTGATCTGTGATAGAAACTGGTAAACATCTAACAATTATGATTGCAGGATT
30	CGTATGTGTCCGTGGAT
	TCGAACAGAAGACAAGAGCACCGCGTCCATTAATGGCAAGGTTGCATTTTCCTGCGAGCAAA
	TGAAGAACAAAACC
	TAAAGATGACTGGAAGAAAGCAAACGCATTGCTTCTCTCTC
	AAGTCCCATTTAGACAT
35	AATAACAGGAATCCAAATAGGCTATTTTCTCTTTCTTTCT
	ACACAAAAAGTTTTT
	TGGGTTATTTATTTTCTCTCTAACAAATTTGTAGCGTTTTGGGTCTTTTTCTGGCTGTCACTAGC
	GTGGCAAATCCAATG
	TCCGCGCACACTTAGGCGCATTGTCAATAAATTCTCCGGCCACCGGAGTGTTACGATCTTTTCC

ATATTTCCGGTAACACATATTCCTTATTCTATGTTGGTTTTGTGTACGGCGTGGGCCTTACTAG
ACAATGATCATCAATA

5

SEQ ID NO:2 DMT amino acid sequence

 ${\tt MQSIMDSSAVNATEATEQNDGSRQDVLEFDLNKTPQQKPSKRKRKFMPKVVVEGKPKRKPRKPA}\\ {\tt ELPKVVVEGKPKRKPR}$

KAATQEKVKSKETGSAKKKNLKESATKKPANVGDMSNKSPEVTLKSCRKALNFDLENPGDARQG

- 10 DSESEIVONSSGANSF
 - ${\tt SEIRDAIGGTNGSFLDSVSQIDKTNGLGAMNQPLEVSMGNQPDKLSTGAKLARDQQPDLLTRNQQ} \\ {\tt CQFPVATQNTQFPME}$
 - NQQAWLQMKNQLIGFPFGNQQPRMTIRNQQPCLAMGNQQPMYLIGTPRPALVSGNQQLGGPQGN KRPIFLNHQTCLPAGN
- 15 QLYGSPTDMHQLVMSTGGQQHGLLIKNQQPGSLIRGQQPCVPLIDQQPATPKGFTHLNQMVATSM SSPGLRPHSQSQVPT
 - TYLHVESVSRILNGTTGTCQRSRAPAYDSLQQDIHQGNKYILSHEISNGNGCKKALPQNSSLPTPIM AKLEEARGSKRQY
 - HRAMGQTEKHDLNLAQQIAQSQDVERHNSSTCVEYLDAAKKTKIQKVVQENLHGMPPEVIEIEDD
- 20 PTDGARKGKNTASIS
 - KGASKGNSSPVKKTAEKEKCIVPKTPAKKGRAGRKKSVPPPAHASEIQLWQPTPPKTPLSRSKPKG KGRKSIQDSGKARG
 - PSGELLCQDSIAEIIYRMQNLYLGDKEREQEQNAMVLYKGDGALVPYESKKRKPRPKVDIDDETTR IWNLLMGKGDEKEG
- 25 DEEKDKKKEKWWEEERRVFRGRADSFIARMHLVQGDRRFSPWKGSVVDSVIGVFLTQNVSDHLS SSAFMSLAARFPPKLS
 - SSREDERNVRSVVVEDPEGCILNLNEIPSWQEKVQHPSDMEVSGVDSGSKEQLRDCSNSGIERFNFL EKSIQNLEEEVLS
 - SODSFDPAIFOSCGRVGSCSCSKSDAEFPTTRCETKTVSGTSQSVQTGSPNLSDEICLQGNERPHLYE
- 30 GSGDVQKQETTN
 - VAQKKPDLEKTMNWKDSVCFGQPRNDTNWQTTPSSSYEQCATRQPHVLDIEDFGMQGEGLGYS WMSISPRVDRVKNKNVP
 - $RRFFRQGGSVPREFTGQIIPSTPHELPGMGLSGSSSAVQEHQDDTQHNQQDEMNKASHLQKTFLDL\\ LNSSEECLTRQSST$
- 35 KQNITDGCLPRDRTAEDVVDPLSNNSSLQNILVESNSSNKEQTAVEYKETNATILREMKGTLADGK KPTSQWDSLRKDVE
 - GNEGRQERNKNNMDSIDYEAIRRASISEISEAIKERGMNNMLAVRIKDFLERIVKDHGGIDLEWLRE SPPDKAKDYLLSI
 - RGLGLKSVECVRLLTLHNLAFPVDTNVGRIAVRMGWVPLQPLPESLQLHLLELYPVLESIQKFLWP
- 40 RLCKLDQRTLYELH

ATAGAAATGATTGAGCA

YQLITFGKVFCTKSRPNCNACPMRGECRHFASAYASARLALPAPEERSLTSATIPVPPESFPPVAIPM
IELPLPLEKSLA
SGAPSNRENCEPIIEEPASPGQECTEITESDIEDAYYNEDPDEIPTIKLNIEQFGMTLREHMERNMELQ
EGDMSKALVAL
HPTTTSIPTPKLKNISRLRTEHQVYELPDSHRLLDGMDKREPDDPSPYLLAIWTPGETANSAQPPEQ
KCGGKASGKMCFD
ETCSECNSLREANSQTVRGTLLIPCRTAMRGSFPLNGTYFQVNELFADHESSLKPIDVPRDWIWDLP
RRTVYFGTSVTSI
FRGLSTEQIQFCFWKGFVCVRGFEQKTRAPRPLMARLHFPASKLKNNKT
SEO ID NO.2 DMT 5' floring acqueres
SEQ ID NO:3 DMT 5' flanking sequence
AAGCTTAAAGCTACCAACATCGAATTTAGTAAAAGACCCATGATTTGAAATTGGAATTGTCGG
CAAAATCGAGAAGATAT
AGAGCCGACACGGGAACAGTGAAAACCACAAAGCGCGTAAGAATGAAACAGTGGGAGAAGG
AAGAGAGAATCTTACCGAT
CATTCGAGGGAAAAGATCAGAGAAAAAATCTGGAAAAAAAGAAATTAAGAGAAAGA
GAGAGAAGAAGTGAGGAG
GAAGATGCAGTGAAGACTGCTATAGCCACATCCCACATGGTGTGATGAGAGAGA
GAGGTTAAAGCAGCAAAT
TGTGGAGAGATAAAGAGAGAGAGAGAGAGAGTCAAGTTCGTCGTCGTCTTTAAAAGAA
AGAATCCTATATTTGCCT
TTTTCTTTACTACTTTTTTCAGACTATTTGCTTATTTTGCCTCAAACTTTTTTGATTGTCACTT
TTCGATCCTAAAGT
GTTTGACAATTTACCTGCCTTTTTCTCCAAGAAAATCAGAACAGACCACAGCAAATTTATGTA
TTTTCTATTAAAAAAG
AAAGAAAGAATTCATATTACTTATAGAATTAAAAGCTAAGCAGTTGAAAACGTGAAAGCAGA
ATTTCTAAAAAAAATAGT
AAACTGCTACAAACTTATTTATGTGTATATAACATATCTATAAAGAAACTCAAATATATGATA
AATCATTTAACAAATAAAAAAAAAAAAAAAAAAAAAAA
TTCTATGAAATAAAAAAAAAAAAAAAAAAAAAAAAAAAA
ACATTTATAATTTCTAAAACTATGGTGTAATTTTGCTGAAATCAAAAAGAAAAGAAAG
TATATCATAAGTTTCAT TATTGTATCAAACTTTCAAATTTCATGTAATTTGAAAGGAAAAAAATTAAGATATAATGTTGTT
TTTGTTTCTTATGTTA CATTTTCATGGGAATATATTCATAACAAAAAATGTATTTTAATATGATGAGAGAGA
CATTTTCATGGAATATATTCATAACAAAAAATGTATTTTAATATGATGAGAGATTACCATCC
AAAAGGTCGAACTTAT

	AACCICAAAAATGICITCITAGGATCACAAAATCITTCCTTAGCTTATTAAAGCCGGGAGTTC
	AACTCTCTCCCTTG
	TAGACTTTTTGTTTTCAAAATCTTTTTCTTTCAAAAAATCAATAATTAGTTAATGGGCATAATATT
	TGGTTTTAATTAAGT
5 '	CCATAGATTTTTTAGGACCATCTCTAATCACGACAAATATCCTAAATTGTAACACATTTAAAAC
	TTAAAAGTATTGCATT
	CACAATCCTTAAAATATATATATATATATATATATATAT
	ATAGAAACGATAACTC
	CTTACTCAACAATTAGCCCAAAAAAACATCCATAATGCATTTAAACTAGGAATTTTAACAAAC
10	TCAAATAGGTTGGTAGT
	TAAAAAAAAAACAAATAGTAGATGTACATACGTACCTTTAAAAAATATATACTCATATCGAAAGT
	TTTAAATTTTGCGAAAT
	TAAATACATTTATCTATCAATTAAAATACATTTAATAATGCATAATTCTGTAATATCTATC
	AATTTCCATATAGAAC
15	CAAAACAAAATAAACATATCAAATAGTTTTAACTTAACAAAAACGTTAGGGAAAAGTTGACCT
	AACTAGCTTGATTGACG
	TTGAACTTGTCAATGCGAAAGCGATATTTCCAATATATACTACATGTAGTATTATTTAT
	AGTTTCTAAAAAGGTG
	TTGAGTGGATTGTTACTTGTTGGAGGATGCTATTTTTTCCTTCTTGCCATAATATTTTACGAGTA
20	TGGGATAACTACATA
	CTCATGATTATGAAACGCTCACTTTATTTGAAAAACCTCCTAATACACCAAATATGTCACTAGA
	TTCCAAAACGTAGACC
	AATTGTATCTAATCTCAAATTCTCAATCAAAGTATTAATTTACCGATGGTAAGAAAAGTTAACC
	GATATAATTATCAAAA
25	GAAAGAATAAGTCAACAGATTCTTAATCTCTTTATTTTGGTATATGAACATTTGTACAAAAATC
	TCAAAAGATATGTAAC
	TGTTTAAAATATAAATTCACTGAGATTAATTCTTCAGACTCGTGTTAGCTATAATAATGTCAAG
	AGTTCTTCTTGTTTCA
	AGGAAAAACCTTAAAGATATGTATATTTTCTGTAATTATGATGATATAATTTGCTATTCATTGT
30	CACAAACATTACTTTA
	AAAAATCGTATTTCATTACTACAATGTTGACTAAGAACAAAAATACATTGATTATTGATATAT
	CGTCAACTGAATTTTC
	TTCCGAGGGATATAATTCTCAAACATAGCAAGAATCTCATAATAATGTTTCGTGACTACCTTTA
	GACGAAATTTTTTAA
35	GATTCGTAACGTGACTTATGGTCTCTTGCTGTGGGGGTCAATGCGAATAAATCTAAATGTATG
	GGAGTCAAATAAAATAC
	CAAGAAAAATAAAGGAGCAGCACCCAATAAACTATATGGGACCAGAAATCCTTTCATTGGTTT
	AAAATAGGATTATCCCG
	AAAGATGAAGGACTAAATTGAAACTGATTGGGGGTAGGAAGAGATCCGTCACAATCATTAAT
40	GGCTTCCACGCGGAAACT

10

15

TGTCGTTTATACAATTTCATTAACTTTCGGGTCGGGTTTATATTCCAAATGGGTCAAATAATAT **TAGTTTAATACACTAA** CGGAGTAATTAATTGGTGACTACAATTTTATCAGTTTGGTGCAATTAGAAACGAACATAGTCG TAAAATACGAGTTCGGT GAATATATGGAAATTA TTAGATACTCTAGCGAAAATAGTGATTATGAGCGTTTTACAAAAATACGATTTTAGCATTGAA CTTCCTTTATGTAATTC GGTCAAATGTTGGCATGAAGAAGCAAGTTTGCAACATTAAATTTCATTTAAAAAATCGTGTTGA CATACTTTAAAATCTAA ATATAGGAAGACCAAAACATTAAATTTAGTAAGATTCTAATGAACATTTATAAGTTATAA CTTATAACCAACAAAG TTGGGTTTAGCGTTGTTGCTTTATCTGAAAACTTGCAAACTAAACCATTTTAATAGGACTAATG ACAATTAACAACAAAA TACACTTAAGCAACACGTCCTCGTGAATATAATTTGGGCCTCAGGCCCATATTGCTAACGCC **AACTGATATTTCACTTT**

20 SEQ ID NO:4 DMT 3' flanking sequence

ATTCCTTCTCACCACCACTCTCTCTCTATCTCTATCTCTAACGGCATAGCTGACTCAGT

SEQ ID NO:5 DMT cDNA sequence

GTTCTCCGGCATTGACTCGCCTGAGAATCAGAAAGCTTAGATCGGTGAGCTTTTAGCTCCATTT
TCTGTTTATTTACATA

- 30 TTATTTCCTTTTTTTCTCCCCTTTTTTATCTGGAATTTGTTCTGCTAAATTTTCCAGCTGTTA CATTTTCCGATCAC
 - GAGAAGAATCACTGGGTTTTTATGTTAATCAATACATGTTCCTGTTTTCTGATCATAAATCTCA GCTATTAACACCTGAT
 - TTTGATTCTGCGTAATAAAAACCTCTGATTTGCTTTTATCTTCACTTTCCCCATAAACATTGCTT
- 35 ACTITATTCGCTCTT

CTTTTACCGTTTCCAGCTAAAAAATTCTTCGCTATTCAATGTGTTTCTCGTTTTGTTGATGAGAA AAATATCTGACAAAA

 ${\tt AATCATTTATTGCATTTTATGGTGCAGATTCTTAGTTAATGTCGCCTTCTCTAACCAAGTCAGATTAAAAAGGAGTGTTC}$



- 35 AGTCACACTCAAAAGTTGCAGAAAAGCTTTGAATTTTGACTTGGAGAATCCTGGAGATGCGAG
 GCAAGGTGACTCTGAGT
 CTGAAATTGTCCAGAACAGTAGTGGCGCAAACTCGTTTTCTGAGATCAGAGATGCCATTGGTG
 GAACTAATGGTAGTTTC
 CTGGATTCAGTGTCACAAATAGACAAGACCAATGGATTGGGGGGCTATGAACCAGCCACTTGAA
 40 GTGTCAATGGGAAACCA

40

AGAGAAGTGGTGGGAAGAA

	GCCAGATAAACTATCTACAGGAGCGAAACTGGCCAGAGACCAACAACCTGATTTATTGACTAG
	AAACCAGCAATGCCAGT
	TCCCAGTGGCAACCCAGAACACCCAGTTCCCAATGGAAAACCAACAAGCTTGGCTTCAGATGA
	AAAACCAACTTATTGGC
5	TTTCCATTTGGTAACCAGCAACCTCGCATGACCATAAGAAACCAGCAGCCTTGCTTG
	GGTAATCAACAACCTAT
	GTATCTGATAGGAACTCCACGGCCTGCATTAGTAAGTGGAAACCAGCAACTAGGAGGTCCCCA
	AGGAAACAAGCGGCCTA
	TATTTTTGAATCACCAGACTTGTTTACCTGCTGGAAATCAGCTATATGGATCACCTACAGACAT
10	GCATCAACTTGTTATG
	TCAACCGGAGGCAACAACATGGACTACTGATAAAAAACCAGCAACCTGGATCATTAATAAG
	AGGCCAGCAGCCTTGCGT
	ACCTTTGATTGACCAGCAACCTGCAACTCCAAAAGGTTTTACTCACTTGAATCAGATGGTAGCT
	ACCAGCATGTCATCGC
15	CTGGGCTTCGACCTCATTCTCAGTCACAAGTTCCTACAACATATCTACATGTGGAATCTGTTTC
	CAGGATTTTGAATGGG
	A CTACAGGTACATGCCAGAGAAGCAGGGCTCCTGCATACGATTCTTTACAGCAAGATATCCAT
	CAAGGAAATAAGTACAT
	ACTTTCTCATGAGATATCCAATGGTAATGGGTGCAAGAAGCGTTACCTCAAAACTCTTCTCTG
20	CCAACTCCAATTATGG
	${\tt CTAAACTTGAGGAAGCCAGGGGCTCGAAGAGACAGTATCATCGTGCAATGGGACAGACGGAA}$
	AAGCATGATCTAAACTTA
	GCTCAACAGATTGCTCAATCACAAGATGTGGAGAGACATAACAGCAGCACGTGTGTGGAATA
	TTTAGATGCTGCAAAGAA .
25	AACGAAAATCCAGAAAGTAGTCCAAGAAAATTTGCATGGCATGCCACCTGAGGTTATAGAAA
	TCGAGGATGATCCAACTG
	ATGGGGCAAGAAAAGGTAAAAATACTGCCAGCATCAGTAAAGGTGCATCTAAAGGAAACTCG
	TCTCCAGTTAAAAAGACA
	GCAGAAAAGGAGAAATGTATTGTCCCAAAAACGCCTGCAAAAAAGGGTCGAGCAGGTAGAAA
30	AAAATCAGTACCTCCGCC
	TGCTCATGCCTCAGAGATCCAGCTTTGGCAACCTACTCCTCCAAAGACACCTTTATCAAGAAG
	CAAGCCTAAAGGAAAAG
	GGAGAAAGTCCATACAAGATTCAGGAAAAGCAAGAGGTCCATCAGGAGAACTTCTGTGTCAG
	GATTCTATTGCGGAAATA
35	ATTTACAGGATGCAAAATCTGTATCTAGGAGACAAAGAAAAGAAAAGAAAAAAAA
	GGTCTTGTACAAAGGAGA
	TGGTGCACTTGTTCCCTATGAGAGCAAGAAGCGAAAACCAAGACCCAAAGTTGACATTGACG
	ATGAAACAACTCGCATAT
	GGAACTTACTGATGGGGAAAGGAGATGAAAAAGAAGGGGGATGAAGAAG

GAAAGAAGAGTCTTCCGAGGAAGGGCTGATTCCTTCATCGCTCGC
GATAGACGTTTTTCGCC
ATGGAAGGGATCGGTGGTTGATTCGGTCATTGGAGTTTTCCTTACACAGAATGTCTCGGATCA
CCTTTCAAGCTCTGCGT
TCATGTCTCTAGCTGCTCGATTCCCTCCAAAATTAAGCAGCAGCCGAGAAGATGAAAGGAATG
TTAGAAGCGTAGTTGTT
GAAGATCCAGAAGGATGCATTCTGAACTTAAATGAAATTCCTTCGTGGCAGGAAAAGGTTCAA
CATCCATCTGACATGGA
AGTTTCTGGGGTTGATAGTGGATCAAAAGAGCAGCTAAGGGACTGTTCAAACTCTGGAATTGA
AAGATTTAATTTCTTAG
AGAAGAGTATTCAAAAATTTAGAAGAGGAAGTATTATCATCACAAGATTCTTTTGATCCGGCGA
TATTTCAGTCGTGTGGG
AGAGTTGGATCCTGTTCATGTTCCAAATCAGACGCAGAGTTTCCTACAACCAGGTGTGAAACA
AAAACTGTCAGTGGAAC
ATCACAATCAGTGCAAACTGGGAGCCCAAACTTGTCTGATGAAATTTGTCTTCAAGGGAATGA
GAGACCGCATCTATATG
AAGGATCTGGTGATGTTCAGAAACAAGAAACTACAAATGTCGCTCAGAAGAAACCTGATCTTC
AAAAACAATGAATTGG
A AAGACTCTGTTTTTGGTCAGCCAAGAAATGATACTAATTGGCAAACAACTCCTTCCAGC
AGCTATGAGCAGTGTGC
GACTCGACAGCCACATGTACTAGACATAGAGGATTTTGGAATGCAAGGTGAAGGCCTTGGTTA
TTCTTGGATGTCCATCT
${\tt CACCAAGAGTTGACAGAGTAAAGAACAAAAATGTACCACGCAGGTTTTTCAGACAAGGTGGA}$
AGTGTTCCAAGAGAATTC
A CAGGTCAGATCATACCATCAACGCCTCATGAATTACCAGGAATGGGATTGTCCGGTTCCTCA
AGCGCCGTCCAAGAACA
${\tt CCAGGACGATACCCAACATAATCAACAAGATGAGATGAATAAAGCATCCCATTTACAAAAAA}$
CATTTTTGGATCTGCTCA
ACTCCTCTGAAGAATGCCTTACAAGACAGTCCAGTACCAAACAGAACATCACGGATGGCTGTC
TACCGAGAGATAGAACT
${\tt GCTGAAGACGTGGTTGATCCGCTCAGTAACAATTCAAGCTTACAGAACATATTGGTCGAATCA}$
AATTCCAGCAATAAAGA
GCAGACGCAGTTGAATACAAGGAGACAAATGCCACTATTTTACGAGAGATGAAAGGGACGC

- 25 CTACAAGCCAGTGGGATAGTCTCAGAAAAGATGTGGAGGGGAATGAAGGGAGACAGGAACG
 AAACAAAAACAATATGGAT
 TCCATAGACTATGAAGCAATAAGACGTGCTAGTATCAGCGAGATTTCTGAGGCTATCAAGGAA
 AGAGGGATGAATAACAT
 GTTGGCCGTACGAATAAGGATTTCCTAGAACGGATAGTTAAAGATCATGGTGGTATCGACCT
- 40 TGAATGGTTGAGAGAAT

TTGCTGATGGGAAAAAGC

TTATTTCATTCATAGA

AAAACTCGAG

CTCCTCCTGATAAAGCCAAGGACTATCTCTTGAGCATAAGAGGTCTGGGTTTGAAAAGTGTTG
AATGCGTGCGACTCTTA
ACACTCCACAATCTTGCTTTCCCTGTTGACACGAATGTTGGAAGGATAGCAGTTAGGATGGGA
TGGGTGCCTCTACAACC
CCTACCTGAATCACTTCAGTTACACCTCCTGGAGCTATACCCAGTGCTCGAGTCCATCCA
TTTCTTTGGCCAAGAC
TTTGCAAACTCGATCAACGAACACTGTATGAATTACACTACCAACTGATTACGTTTGGAAAGG
TATTTTGCACAAAGAGT
AGACCAAATTGTAATGCATGTCCAATGAGAGGAGAGTGCAGACACTTTGCCAGTGCTTATGCT
AGTGCAAGACTTGCTTT
ACCGCACCAGAGGAGGAGCTTAACAAGTGCAACTATTCCGGTCCCTCCC
TCCTGTAGCCATCCGA
TGATAGAACTACCTCTTCCGTTGGAGAAATCCCTAGCAAGTGGAGCACCATCGAATAGAGAA
ACTGTGAACCAATAATT
GAAGAGCCGGCCTCGCCCGGGCAAGAGTGCACTGAAATAACCGAGAGTGATATTGAAGATGC
TTACTACAATGAGGACCC TGACGAGATCCCAACATAAAACTCAACATTGAACAGTTTGGAATGACTCTACGGGAACACAT
GGAAAGAACATGGAGC
TCCAAGAAGTGACATGTCCAAGGCTTTGGTTGCTTTGCATCCAACAACTACTTCTATTCCAAC
TCCCAAACTAAAGAAC
ATTAGCCGTCTCAGGACAGCACCAAGTGTACGAGCTCCCAGATTCACATCGTCTCCTTGAT
GGTATGGATAAAAGAGA
ACCAGATGATCCAAGTCCTTATCTCTTAGCTATATGGACACCAGGTGAAACAGCGAATTCGGC
ACAACCGCCTGAACAGA
AGTGTGGAGGGAAAGCGTCTGGCAAAATGTGCTTTGACGAGACTTGTTCTGAGTGTAACAGTC
TGAGGGAAGCAAACTCA
CAGACAGTTCGAGGAACTCTTCTGATACCTTGTCGGACTGCCATGAGAGGAAGTTTTCCGCTC
AACGGGACATATTTCCA
AGTCAACGAGTTATTTGCAGACCACGAGTCCAGTCTCAAACCCATCGATGTTCCTAGAGATTG
GATATGGGATCTCCCAA
GAAGGACTGTTTACTTCGGAACATCAGTAACATCAATATTCAGAGGTCTTTCAACGGAGCAGA
TACAGTTCTGCTTTTGG
AAAGGATTCGTATGTGTCCGTGGATTCGAACAGAAGACAAGAGCACCGCGTCCATTAATGGCA
AGGTTGCATTTTCCTGC
GAGCAAATTGAAGAACAAAAACCTAAAGATGACTGGAAGAAAGCAAACGCATTGCTTCTC
TGCTCTCTCTATTTAAA

	SEQ ID NO:6 5' untranslated region of DMT
	GTTCTCCGGCATTGACTCGCCTGAGAATCAGAAAGCTTAGATCGGTGAGCTTTTAGCTCCATTT
	TCTGTTTATTTACATA
5	TTATTTCCTTTTTTCTCTCCCTTTTTTATCTGGAATTTGTTCTGCTAAATTTTCCAGCTGTTA
	CATTITCCGATCAC
	GAGAAGAATCACTGGGTTTTTATGTTAATCAATACATGTTCCTGTTTTCTGATCATAAATCTCA
	GCTATTAACACCTGAT
	TTTGATTCTGCGTAATAAAAACCTCTGATTTGCTTTTATCTTCACTTTCCCCATAAACATTGCTT
10	ACTTTATTCGCTCTT
	$\tt CTTTTACCGTTTCCAGCTAAAAAATTCTTCGCTATTCAATGTGTTTCTCGTTTTGTTGATGAGAA$
	AAATATCTGACAAAA
	AATCATTTATTGCATTTTATGGTGCAGATTCTTAGTTAATGTCGCCTTCTCTAACCAAGTCAGA
	TTAAAAAGGAGTGTTC
15	$\tt GTCCATGTTGCTTTTGGTGTTTTGGAGAGAGTTTTCGGAGAGTTAGGTGAGTGTTATTTGG$
	GGTGAGGTAGTGATAA
	${\tt GGTTTGAAGGGGGAGTGATTCATCAAGTGTGTTATGAATTCGAGGGCTGATCCGGGGGATAGA}$
	TATTTTCGAGTTCCTTT
	GGAGAATCAAACTCAACAAGAGTTCATGGGTTCTTGGATTCCATTTACACCCAAAAAACCTAG
20	ATCAAGTCTGATGGTAG
	ATGAGAGAGTGATAAACCAGGATCTAAATGGGTTTCCAGGTGGTGAATTTGTAGACAGGGGA
	TTCTGCAACACTGGTGTG
	GATCATAATGGGGTTTTTGATCATGGTGCTCATCAGGGCGTTACCAACTTAAGTATGATGATCA
	ATAGCTTAGCGGGATC
25	ACATGCACAAGCTTGGAGTAATAGTGAGAGAGATCTTTTTGGGCAGGAGTGAGGTGACTTCTCC
	TTTAGCACCAGTTATCA
	GAAACACCACCGGTAATGTAGAGCCGGTCAATGGAAATTTTACTTCAGATGTGGGTATGGTAA
	ATGGTCCTTTCACCCAG
	AGTGGCACTTCTCAAGCTGGCTATAATGAGTTTGAATTGGATGACTTGTTGAATCCTGATCAGA
30	TGCCCTTCTCCTTCAC
	AAGCTTGCTGAGTGGTGGGGATAGCTTATTCAAGGTTCGTCAATGTGAGTGA
	TCAGTTTTTTTTC
	CCTTTCTTCCGTTCTTGCAGTACTTAGAGTAGAACATGAATTAGAATATCTTAAGAAAGTCATG
	GTTTTGAACAGATGGA
35	CCTCCAGCGTGTAACAAGCCTCTTTACAATTTGAATTCACCAATTAGAAGAGAAGCAGTTGGG
	TCAGTCTGTGAAAGTTC
	GTTTCAATATGTACCGTCAACGCCCAGTCTGTTCAGAACAGGTGAAAAGACTGGATTCCTTGA
	۸ C A G A T A GTT A C A A CT A

CTGGACATGAAATCCCAGAGCCGAAATCTGACAAAAGT


SEQ ID NO:7 >Arabidopsis thaliana DMT1(1DMT5) gene sequence from BAC T32M21 (gi 7406444);

	64441	tcactagatt	ccaaaacgta	gaccaattgt	atctaatctc	aaattctcaa	tcaaagtatt
	64501	aatttaccga	tggtaagaaa	agttaaccga	tataattatc	aaaagaaaga	ataagtcaac
5	64561	agattcttaa	tctctttatt	ttggtatatg	aacatttgta	caaaaatctc	aaaagatatg
	64621	taactgttta	aaatataaat	tcactgagat	taattcttca	gactcgtgtt	agctataata
	64681	atgtcaagag	ttcttcttgt	ttcaaggaaa	aaccttaaag	atatgtatat	tttctgtaat
	64741	tatgatgata	taatttgcta	ttcattgtca	caaacattac	tttaaaaaat	cgtattttca
	64801	ttactacaat	gttgactaag	aacaaaaata	cattgattat	tgatatatcg	tcaactgaat
10	64861	tttcttccga	gggatataat	tctcaaacat	agcaagaatc	tcataataat	gtttcgtgac
	64921	tacctttaga	cgaaattttt	ttaagattcg	taacgtgact	tatggtctct	tgctgtgggg
	64981	gtcaatgcga	ataaatctaa	atgtatggga	gtcaaataaa	ataccaagaa	aaataaagga
	65041	gcagcaccca	ataaactata	tgggaccaga	aatcctttca	ttggtttaaa	ataggattat
	65101	cccgaaagat	gaaggactaa	attgaaactg	attgggggta	ggaagagatc	cgtcacaatc
15	65161	attaatggct	tccacgcgga	aacttgtcgt	ttatacaatt	tcattaactt	tcgggtcggg
	65221	tttatattcc	aaatgggtca	aataatatta	gtttaataca	ctaacggagt	aattaattgg
	65281	tgactacaat	tttatcagtt	tggtgcaatt	agaaacgaac	atagtcgtaa	aatacgagtt
	65341	cggtgttata	cctttattta	cgttaaaaaa	atacgagaat	tttgtgtcaa	atttcaaatt
		aatttcatga					
20		ttacaaaaat	•				
		gaagaagcaa	_	_	_		
		ctaaatatag	_				
		gttataactt					
	65701	aaactaaacc	attttaatag	gactaatgac	aattaacaac	aaaatacact	taagcaacaa
25		cgtcctcgtg	_				_
		ctttattcct					
		gactcagtgt					
		ttagctccat					
		ctggaatttg	-				
30 -		tgggttttta	_		_		
		cacctgattt					
		ataaacattg					
		tattcaatgt				-	
	66301	cattttatgg	tgcagattct	tagttaatgt	cgccttctct	aaccaagtca	gattaaaaag
35		gagtgttcgt					
		agtgttattt					
		t atg aattcg					
		tcaacaagag					
		gatggtagat					
40		agacagggga					
		tcagggcgtt	-				•
		ttggagtaat					
		agttatcaga					
		gggtatggta	_				
45		tgaattggat					
		tggtggggat					
		ttttttccc					
		aagaaagtca					_
		gaattcacca					
50		accgtcaacg					_
, -		tacaactact					

67321 tacaactact ggacatgaaa tcccagagcc gaaatctgac aaaagtATGc agagcattat

	67381	ggactcgtct	gctgttaatg	cgacggaagc	tactgaacaa	aatgatggca	gcagacaaga
	67441	tgttctggag	ttcgacctta	acaaaactcc	tcagcagaaa	ccctccaaaa	ggaaaaggaa
	67501	gttcatgccc	aaggtggtcg	tggaaggcaa	acctaaaaga	aagccacgca	aacctgcaga
_	67561	acttcccaaa	gtggtcgtgg	aaggcaaacc	taaaaggaag	ccacgcaaag	ctgcaactca
5	67621	ggaaaaagtg	aaatctaaag	aaaccgggag	tgccaaaaag	aaaaatttga	aagaatcagc
	67681	aactaaaaag	ccagccaatg	ttggagatat	gagcaacaaa	agccctgaag	tcacactcaa
	67741	aagttgcaga	aaagctttga	attttgactt	ggagaateet	ggagatgcga	ggcaaggtga
	67801	ctctgagtct	gaaattgtcc	agaacagtag	tggcgcaaac	tegttttetg	agatcagaga
10	67861	tgccattggt	ggaactaatg	gtagtttcct	ggattcagtg	tcacaaatag	acaagaccaa
10	67921	tggattgggg	gctatgaacc	agccacttga	agtgtcaatg	ggaaaccagc	cagataaact
	67981	atctacagga	gcgaaactgg	ccagagacca	acaacctgat	ttattgacta	gaaaccagca
	68041	atgccagttc	ccagtggcaa	cccagaacac	ccagttccca	atggaaaacc	aacaagcttg
	.68101	gcttcagatg	aaaaaccaac	ttattggctt	tccatttggt	aaccagcaac	ctcgcatgac
	68161	cataagaaac	cagcagcctt	gcttggccat	gggtaatcaa	caacctatgt	atctgatagg
15	68221	aactccacgg	cctgcattag	taagtggaaa	ccagcaacta	ggaggtcccc	aaggaaacaa
	68281	gcggcctata	tttttgaatc	accagacttg	tttacctgct	ggaaatcagc	tatatggatc
	68341	acctacagac	atgcatcaac	ttgttatgtc	aaccggaggg	caacaacatg	gactactgat
	68401	aaaaaaccag	caacctggat	cattaataag	aggccagcag	ccttgcgtac	ctttgattga
	68461	ccagcaacct	gcaactccaa	aaggttttac	tcacttgaat	cagatggtag	ctaccagcat
20	68521	gtcatcgcct	gggcttcgac	ctcattctca	gtcacaagtt	cctacaacat	atctacatgt
	68581	ggaatctgtt	tccaggattt	tgaatgggac	tacaggtaca	tgccagagaa	gcagggctcc
	68641	tgcatacgat	tctttacagc	aagatatcca	tcaaggaaat	aagtacatac	tttctcatga
	68701	gatatccaat	ggtaatgggt	gcaagaaagc	gttacctcaa	aactcttctc	tgccaactcc
	68761	aattatggct	aaacttgagg	aagccagggg	ctcgaagaga	cagtatcatc	gtgcaatggg
25	68821	acagacggaa	aagcatgatc	taaacttagc	tcaacagatt	gctcaatcac	aagatgtgga
	68881	gagacataac	agcagcacgt	gtgtggaata	tttagatgct	gcaaagaaaa	cgaaaatcca
	68941	gaaagtagtc	caagaaaatt	tgcatggcat	gccacctgag	gttatagaaa	tcgaggatga
	69001	tccaactgat	ggggcaagaa	aaggtaaaaa	tactgccagc	atcagtaaag	gtgcatctaa
_ :	69061	aggaaactcg	tctccagtta	aaaagacagc	agaaaaggag	aaatgtattg	tcccaaaaac
30_	69121	gcctgcaaaa	aagggtcgag	caggtagaaa	aaaatcagta	cctccgcctg	ctcatgcctc
	69181	agagatccag	ctttggcaac	ctactcctcc	aaagacacct	ttatcaagaa	gcaagcctaa
	69241	aggaaaaggg	agaaagtcca	tacaagattc	aggaaaagca	agaggtaact	aatgtattct
	69301	acaatctctg	tgatataatt	ttgagatttt	agtaactgat	gtgtccaaac	cagctcctta
	69361	tcactgttgg	tgcgttgtat	aggtccatca	ggagaacttc	tgtgtcagga	ttctattgcg
35	69421	gaaataattt	acaggatgca	aaatctgtat	ctaggagaca	aagaaagaga	acaagagcaa
	69481	aatgcaatgg	tcttgtacaa	aggagatggt	gcacttgttc	cctatgagag	caagaagcga
	69541	aaaccaagac	ccaaagttga	cattgacgat	gaaacaactc	gcatatggaa	cttactgatg
	69601	gggaaaggag	atgaaaaaga	aggggatgaa	gagaaggata	aaaagaaaga	gaagtggtgg
	69661	gaagaagaaa	gaagagtett	ccgaggaagg	gctgattcct	tcatcgctcg	catgcacctg
40	69721	gtacaaggtg	aagatccact	tctcttctca	actccatttt	tattcacaca	aattagtaga
	69781	atactcaaaa	atgatgtttt	gtttgcaaaa	ttttaaaatt	cactagttaa	ccatgtcaaa
	69841	taatattcat	aatgcatctt	gtgaagaaca	ggtgtgcatt	tatggtgaca	gctgaatggt
	69901	ttatgtgcct	attatttctt	ttactgctat	agatgaccaa	ttgaacttaa	acgtttacag
4.5	69961	gagatagacg	tttttcgcca	tggaagggat	cggtggttga	ttcggtcatt	ggagttttcc
45	70021	ttacacagaa	tgtctcggat	cacctttcaa	ggtatatgag	ttgccttaat	aaattgagtt
	70081	ccaaaacata	gaaattaacc	catggtggtt	ttacaatgca	gctctgcgtt	catgtctcta
	70141	gctgctcgat	tccctccaaa	attaagcagc	agccgagaag	atgaaaggaa	tgttagaagc
	70201	gtagttgttg	aagatccaga	aggatgcatt	ctgaacttaa	atgaaattcc	ttcgtggcag
.	70261	gaaaaggttc	aacatccatc	tgacatggaa	gtttctgggg	ttgatagtgg	atcaaaagag
50	70321	cagctaaggg	actgttcaaa	ctctggaatt	gaaagattta	atttcttaga	gaagagtatt
	70381	caaaatttag	aagaggaagt	attatcatca	caagattctt	ttgatccggc	gatatttcag

	70441	tcgtgtggga	gagttggatc	ctgttcatgt	tccaaatcag	acgcagagtt	tcctacaacc
	70501	aggtgtgaaa	caaaaactgt	cagtggaaca	tcacaatcag	tgcaaactgg	gagcccaaac
	70561	ttgtctgatg	aaatttgtct	tcaagggaat	gagagaccgc	atctatatga	aggatctggt
-	70621	gatgttcaga	aacaagaaac	tacaaatgtc	gctcagaaga	aacctgatct	tgaaaaaaca
5	70681	atgaattgga	aagactctgt	ctgttttggt	cagccaagaa	atgatactaa	ttggcaaaca
	70741	actccttcca	gcagctatga	gcagtgtgcg	actcgacagc	cacatgtact	agacatagag
	70801	gattttggaa	tgcaaggtga	aggccttggt	tattcttgga	tgtccatctc	accaagagtt
		gacagagtaa					
10	70921	agagaattca	caggtcagat	cataccatca	acgcctcatg	aattaccagg	aatgggattg
10	70981	teeggtteet	caagcgccgt	ccaagaacac	caggacgata	cccaacataa	tcaacaagat
	71041	gagatgaata	aagcatccca	tttacaaaaa	acatttttgg	atctgctcaa	ctcctctgaa
	71101	gaatgcctta	caagacagtc	cagtaccaaa	cagaacatca	cggatggctg	tctaccgaga
	71161	gatagaactg	ctgaagacgt	ggttgatccg	ctcagtaaca	attcaagctt	acagaacata
	71221	ttggtcgaat	caaattccag	caataaagag	cagacggcag	ttgaatacaa	ggagacaaat
15	71281	gccactattt	tacgagagat	gaaagggacg	cttgctgatg	ggaaaaagcc	tacaagccag
	71341	tgggatagtc	tcagaaaaga	tgtggagggg	aatgaaggga	gacaggaacg	aaacaaaac
	71401	aatatggatt	ccatagacta	tgaagcaata	agacgtgcta	gtatcagcga	gatttctgag
	71461	gctatcaagg	aaagagggat	gaataacatg	ttggccgtac	gaattaaggt	aaatctacta
• •	71521	atttcagttg	agaccctcat	caaatctgtc	agaaggettg	aacatcagta	aattatgtaa
20	71581	ccatatttac	aacattgcag	gatttcctag	aacggatagt	taaagatcat	ggtggtatcg
	71641	accttgaatg	gttgagagaa	tetectectg	ataaagccaa	gtgggtaaat	cacatttta
	71701	gtgactgcaa	cactagcacg	atcgatttac	tcaacaatta	cgtcaaactg	agtattaaca
	71761	agttgctcat	gaacatttca	cagggactat	ctcttgagca	taagaggtct	gggtttgaaa
0.5	71821	agtgttgaat	gcgtgcgact	cttaacactc	cacaatcttg	ctttccctgt	gagtcagact
25	71881	attccattat	ctactaaaaa	cttagaataa	ctccggctaa	ctaagctgga	acttgtattg
	71941	atgatatgaa	ggttgacacg	aatgttggaa	ggatagcagt	taggatggga	tgggtgcctc
	72001	tacaacccct	acctgaatca	cttcagttac	acctcctgga	gctgtaagtt	tctttttgtt
	72061	tgtcatctaa	acaacgaaat	ttttatgcaa	gtcataacca	tgctgtgttt	tcacagatac
20		ccagtgctcg					
30		acactgtatg					
	72241	agaattctat	tgtcattact	cattactaac	agcgaaatta	attaacgttc	tttttcttac
		tcaggtatga					
		tgaatttatc					
25		catgtgaatt					
35		atgagaggag					
		cttatatgca	_				
		tcagtttcat					
		gaatcatgga	_		_		-
40		tgccaaagta					
40		acttgcttta					
		cgagtcctat			_	_	
		cctagcaagt				_	
		ctcgcccggg		_			
45		tgaggaccct					-
43		acgggaacac					
		tgctttgcat				_	
		caggacagag					
		tttggggaaa	_				
50		catttttata				_	_
50		gcattgaaca				_	
	/3441	aaatgacaag	ycaactgata	aatgugtegt	ccatttgcag	ycacgagete	ccagattcac



SEQ ID NO:8 ARABIDOPSIS THALIANA DMT2

>DMT2(1DMT2);

5

10

15

20

25

MEKQRREESSFQQPPWIPQTPMKPFSPICPYTVEDQYHSSQLEERRFVGNKDMSGLDHLS FGDLLALANTASLIFSGQTPIPTRNTEVMQKGTEEVESLSSVSNNVAEQILKTPEKPKRK KHRPKVRREAKPKREPKPRAPRKSVVTDGQESKTPKRKYVRKKVEVSKDQDATPVESSAA VETSTRPKRLCRRVLDFEAENGENQTNGDIREAGEMESALQEKQLDSGNQELKDCLLSAP STPKRKRSQGKRKGVQPKKNGSNLEEVDISMAQAAKRRQGPTCCDMNLSGIQYDEQCDYQ KMHWLYSPNLQQGGMRYDAICSKVFSGQQHNYVSAFHATCYSSTSQLSANRVLTVEERRE GIFOGROESELNVLSDKIDTPIKKKTTGHARFRNLSSMNKLVEVPEHLTSGYCSKPQQNN KILVDTRVTVSKKKPTKSEKSQTKQKNLLPNLCRFPPSFTGLSPDELWKRRNSIETISEL LRLLDINREHSETALVPYTMNSQIVLFGGGAGAIVPVTPVKKPRPRPKVDLDDETDRVWK $\verb|LLENINSEGVDGSDEQKAKWWEEERNVFRGRADSFIARMHLVQGDRRFTPWKGSVVDSV|\\$ VGVFLTONVSDHLSSSAFMSLASOFPVPFVPSSNFDAGTSSMPSIOITYLDSEETMSSPP DHNHSSVTLKNTOPDEEKDYVPSNETSRSSSEIAISAHESVDKTTDSKEYVDSDRKGSSV EVDKTDEKCRVLNLFPSEDSALTCQHSMVSDAPQNTERAGSSSEIDLEGEYRTSFMKLLQ GVQVSLEDSNQVSPNMSPGDCSSEIKGFQSMKEPTKSSVDSSEPGCCSQQDGDVLSCQKP TLKEKGKKVLKEEKKAFDWDCLRREAQARAGIREKTRSTMDTVDWKAIRAADVKEVAETI KSRGMNHKLAERIQYLTLNMKIMQGFLDRLVNDHGSIDLEWLRDVPPDKAKEYLLSFNGL GLKSVECVRLLTLHHLAFPVDTNVGRIAVRLGWVPLQPLPESLQLHLLEMYPMLESIQKY LWPRLCKLDOKTLYELHYOMITFGKVFCTKSKPNCNACPMKGECRHFASAFARKFSNIHL FYSARLALPSTEKGMGTPDKNPLPLHLPEPFOREOGSEVVOHSEPAKKVTCCEPIIEEPA SPEPETAEVSIADIEEAFFEDPEEIPTIRLNMDAFTSNLKKIMEHNKELQDGNMSSALVA LTAETASLPMPKLKNISQLRTEHRVYELPDEHPLLAQLEKREPDDPCSYLLAIWTPGETA DSIOPSVSTCIFOANGMLCDEETCFSCNSIKETRSQIVRGTILIPCRTAMRGSFPLNGTY FOVNEVFADHASSLNPINVPRELIWELPRRTVYFGTSVPTIFKGLSTEKIOACFWKGYVC VRGFDRKTRGPKPLIARLHFPASKLKGQQANLA

SEO ID NO:9

30 >DMT2(1DMT2) novel 480 amino acid amino terminus;
MEKQRREESSFQQPPWIPQTPMKPFSPICPYTVEDQYHSSQLEERRFVGNKDMSGLDHLS
FGDLLALANTASLIFSGQTPIPTRNTEVMQKGTEEVESLSSVSNNVAEQILKTPEKPKRK
KHRPKVRREAKPKREPKPRAPRKSVVTDGQESKTPKRKYVRKKVEVSKDQDATPVESSAA
VETSTRPKRLCRRVLDFEAENGENQTNGDIREAGEMESALQEKQLDSGNQELKDCLLSAP
STPKRKRSQGKRKGVQPKKNGSNLEEVDISMAQAAKRRQGPTCCDMNLSGIQYDEQCDYQ

KMHWLYSPNLQQGGMRYDAICSKVFSGQQHNYVSAFHATCYSSTSQLSANRVLTVEERRE GIFQGRQESELNVLSDKIDTPIKKKTTGHARFRNLSSMNKLVEVPEHLTSGYCSKPQQNN KILVDTRVTVSKKKPTKSEKSOTKOKNLLPNLCRFPPSFTGLSPDELWKRRNSIETISEL

5 SEQ ID NO:10

>DMT2 (1DMT2) Nucleotide sequence from BAC F1011 (gi 6598632); 60001 tegetgagee tgggtttett categgaeet ggatetetgg atetateaaa eggtetaega 60061 ggattctcca ttccaaagaa ctatacaata caagaggtac gcaaataatg ccctaaatta 60121 aacctaatcg gcaaaaatcg attgcagtga caacaaatcc tcgttagagg ggaattcaga 10 60181 gcattacaac aatcagtaac cctaagttac aatctaaaaa ttgagatgca taacgcgatt 60241 ctgcgaagaa gacggagaag atagaaggaa tgcttcgaat tcggcaaaaa tgtcagagag 60301 tttggacaat ctccgatcaa ttagggttgt gaattgggga ttttatggag acgagacaaa 60361 aaaaagttga agatcggagc tggttccaaa aatatttagg cccatttaat gacccacatt 60421 ccatgtataa taggcccatc atctaatatt tgacaacaat agaattcttt ggtccggttg 15 60481 aactatctga tttaaaccaa gttaagtgag atcctccaca tatcgaacca gatcttgatt 60541 caggtaacca aaagctaacc gtaaattcag atataaacca aacgaaggga acagagagtt 60601 tacacageta egggtetgtt ttttgtgaca agtgtttgat acaaatttaa gacgaaacta 60661 aaatgggatt tagaaacctt gtacaactct aggactgtta actttacgtt ttcactttct 60721 tacattaact agattggaac agtgtgctct ctcactctta accataagct tgtatttgtt 20 60781 tgcttgccaa cggaTTAggc gaggttagct tgttgtccct tcagtttgct cgccgggaag 60841 tgcaatcttg caatcaaagg cttcggtccc ctcgtctttc gatcaaatcc acgtacacat 60901 acgtacccta cataatatca aaagataagt tatgtttcag aacaagaaga aactgcttaa 60961 tacaaaatgt acctttccaa aagcaagcct gtatcttctc agttgataaa cctgagaaaa 61021 atagagetea agtggttaga acaaetttet tttatataaa caategeate acaateeaat 25 61081 aaagaaaatc ttataccttt gaatatcgta ggaacagagg taccaaaata gaccgttctt 61141 cgaggtaatt cccatatcaa ttcccttggg acattgattg ggtttaggct ggatgcatga 61201 tccgcaaaca cctgtatcaa tagaatacat cacaagtttc aatgcaaata attaaaatga 61261 aagagttgga gttattggag ttcaagtctt acctcattta cttgaaagta cgttccattt 61321 agaggaaaac tacccctcat cgctgttcta caaggaatct gtacaattta caacatatta 30 61381 atctgtagaa aacataagtg tagtaagccg cataaggaga ttgatgcaac tacttaccaa 61441 aattgtccct ctcacaattt gagatctagt ctccttgatg ctgttgcagg agaaacaagt 61501 ctcctcgtca caaagcatac catttgcttg gaatatgcac gtactaacag acggttgaat 61561 agaatcagcc gtctcacctg ttgaataaca catcgattaa agataccgat ttgatttcat 61621 gattaaaaga tatgcaaatc attaaattac ctggcgtcca tatagcaagc aaataagaac 35 61681 atggatcatc aggttctctc ttttccaact gccacaagaa atcacaaaca gctagtcaga 61741 ttttacaata tagacagcac tctatacggc atgtgtcctt atccagttag ctcacatacc 61801 tgagctagaa gaggatgctc gtctggaagt tcgtaactgc aagatacggg aaaagaaaca 61861 agttatggca tagcctgtaa ttattgggaa gtttgtctgc tttccaactt acgagttcat 61921 gcttqqtcaa tcacttaaat attctactct qttcaagctt taataatttt gaaaaatgtg 40 61981 tttctgattt catttttaac ctaagaacga agaaaaacag agaaaaatgg attcttacac 62041 tcggtgttct gtccttaact ggctgatatt cttgagctta ggcattggaa gagaagcagt 62101 ttcagcagta agtgcaacta aagcgctgga catgtttccg tcttgaagtt ccttgttgtg 62161 ttccattatc ttcttcaagt tactggtaaa tgcatccatg tttagcctga tggtaggaat 62221 ttcttctgga tcctcaaaaa acgcctcctc tatgtcagct attgatactt ctgcggtttc 45 62281 tggctccggt gaagcaggct cttcgatgat tggttcacaa catgtgacct tttttgctgg 62341 ttctgagtgc tgtactactt cagaccettg ctctctctgg aatggctctg gcaggtgtag 62401 aggcaaaggg tttttatcag gtgtccccat acctttctct gtacttggta aagcaagcct 62461 tgcactgtaa aacaaatgaa tgttactaaa ttttctgtaa tgatgattca gagcttcgtt

62521 tagatacaga ccaattetea tttaaetggg ttatatttta acaaggaett teeteataga

62581	gtcatagtgg	tactaaaggt	ttaagagaac	atgttgtagc	accttgcaaa	cgcactggca
62641	aaatgtctgc	attctccttt	catcggacat	gcattgcaat	taggtttgct	ctttgtgcaa
62701	aagacctgat	acaataatca	agcagattac	aaacctcatc	atgtgagctg	attttgacat
62761	acgtatatat	gtatttcttt	aatacatacc	tttccaaaag	taatcatctg	gtagtgcaac
62821	tcatacctgt	gagataatag	ggtattaaac	taatgaataa	gtgtattaga	ctgaggcatg
62881	aaaaaaaaa	agttagtgat	aaacatcatt	cttacaatgt	tttttggtcg	agtttgcaga
62941	gacggggcca	aagatacttt	tgaatagatt	caagcatagg	atacctagac	aaaccaaacc
63001	tcagatgtat	taagtaacaa	attacaattt	ccaagtagga	ccattttgaa	aagtgcttac
63061	atttccagaa	gatgcaactg	aagtgactct	gggagcggct	gaaggggcac	ccatccaagt
63121	ctgacggcta	tgcgcccaac	atttgtatca	acctgtcaat	aaattaagtt	catgcatcat
63181	ataattcact	ttttataggg	acagaaacaa	aagtttgatc	cttgcttact	ggaaaggcaa
63241	gatggtgaag	tgttagaagc	cgcacacact	ccacactttt	cagtcccaat	ccgttaaagc
63301	tcagaagata	ttctctgcag	ggttttgtaa	tatacgagag	tacataattc	attattaagt
63361	cactaaaact	gccaaagtag	taatctttgt	ataggttaat	aaagaagaaa	taaaatgctt
63421	cgtctttcaa	acttactttg	ctttatctgg	tggaacatct	ctcaaccatt	caagatcgat
63481	acttccatgg	tcatttacca	gtcgatcaag	gaagccctgc	atgattttca	tgttcagagt
63541	caaatactta	aaatgaatgt	tatcacgaaa	tttagccact	aaattttac	ctgtatacgt
63601	tctgcaagtt	tatggttcat	cccgcgactc	ttgattgttt	cagcaacttc	cttaacatct
63661	gctgctcgta	ttgccttcca	atccacggtg	tccattgtac	ttcttgtttt	ttctctaatt
63721	cctgctctag	cttgggcttc	tcttcttaaa	caatcccagt	caaacgcttt	tttttcctcc
63781	ttcaaaacct	ttttcccttt	ttcttttaag	gtaggtttct	gacaactcaa	aacatcccca
63841	tcttgctgag	agcaacaacc	aggttcacta	ctatcaacag	aggattttgt	gggctctttc
63901	attgactgga	aacccttaat	ttctgagcta	caatcacccg	gagacatatt	tggtgatact
63961	tgattggaat	cttctagaga	gacttgtacc	ccctgtagga	gcttcataaa	ggaagtacga
64021	tactctcctt	ctaagtcgat	ctctgagctt	gatcctgctc	tctctgtatt	ttgaggagca
64081	tcagacacca	tcgaatgttg	acatgtaagt	gcagaatctt	cagatggaaa	caggttcagg
64141	acacgacact	tctcatccgt	cttatcaacc	tctacacttg	agccttttcg	atctgaatca
64201	acatactcct	ttgaatccgt	ggttttgtca	actgattcat	gggctgagat	ggcaatctca
64261	ctactgcttc	tggaggtttc	attgctaggt	acataatcct	tctcctcatc	aggctgtgta
64321	tttttcaaag	taacagaact	gtgattgtga	tcgggtgggc	ttgacatcgt	ttcctctgag
64381	tccaagtacg	ttatttgaat	agaaggcatc	gagcttgttc	cagcgtcaaa	gttactgctc
64441	ggtacaaaag	ggacagggaa	ctgggaagcc	aacgacatga	aagccgaact	acaaggagta
64501	aaaaacatca	aagcaagtta	gttttgtgac	tttttgctgt	cttggattta	gtttgacata
64561	gaattatgta	agagcttgta	ccttgagaga	tggtctgaaa	cattttgagt	gagaaatact
64621	ccaacaacag	aatccacgac	ggatcccttc	caaggcgtaa	aacgtcgatc	ccctgttaga
64681	aaccaaagac	cataacaaga	agcagtagct	gagacatact	aattgaaacc	atgtggttag
64741	aacagaaaca	cataaaagga	caagtgtggt	gtataacctt	gtacaaggtg	catccttgca
64801	ataaatgagt	cagctcgtcc	tcgaaacaca	ttacgttctt	cctcccacca	tttcgccttc
64861	tgctcgtctg	atccgtcaac	accttcgcta	ttaatattct	ccaatagcag	tttccacact
64921	ctgtctgtct	catcgtctag	atcaaccttt	ggtcgtgggc	gtggttttt	aacaggagtt
					tctggctatt	
65041	ggaacgagag	cagtttcaga	atgctccctg	ttgatgtcta	atagacgcaa	tagctcactg
					gagaaagacc	
			_		ttcaaaaaaa	
					aacggcaaag	
					gcttcttttt	
					tgctacaata	
					aattccggaa	
					agagaacatt	
*					cggttaggac	
					cagaaacata	_
		_				

68641 tgaattaaac tataaataga aataaccaaa catataacca caaaagaaga ctatttatat

15

20

25

30

35

68701 aaatatatga gttggaagte atttttggae tattatataa gatetaatta teacaegaeg 68761 tgtggatgta tggttageag agttgtgtte agagagtteg ataaageeat caeteeaaae 68821 atacaaaata teeatacatt gateeaceaa tataaeegge tgtgtgeeaa geaaagtgaa

5 SEQ ID NO:11

ARABIDOPSIS THALIANA DMT3

>DMT3(1DMT3);

MEVEGEVREKEARVKGRQPETEVLHGLPQEQSIFNNMQHNHQPDSDRRRLSLENLPGLYN MSCTOLLALANATVATGSSIGASSSSLSSOHPTDSWINSWKMDSNPWTLSKMQKQQYDVS TPQKFLCDLNLTPEELVSTSTQRTEPESPQITLKTPGKSLSETDHEPHDRIKKSVLGTGS PAAVKKRKIARNDEKSOLETPTLKRKKIRPKVVREGKTKKASSKAGIKKSSIAATATKTS EESNYVRPKRLTRRSIRFDFDLOEEDEEFCGIDFTSAGHVEGSSGEENLTDTTLGMFGHV PKGRRGORRSNGFKKTDNDCLSSMLSLVNTGPGSFMESEEDRPSDSQISLGRQRSIMATR PRNFRSLKKLLORIIPSKRDRKGCKLPRGLPKLTVASKLQLKVFRKKRSQRNRVASQFNA RILDLOWRRONPTGTSLADIWERSLTIDAITKLFEELDINKEGLCLPHNRETALILYKKS YEEOKAIVKYSKKOKPKVOLDPETSRVWKLLMSSIDCDGVDGSDEEKRKWWEEERNMFHG RANSFIARMRVVOGNRTFSPWKGSVVDSVVGVFLTONVADHSSSSAYMDLAAEFPVEWNF NKGSCHEEWGSSVTQETILNLDPRTGVSTPRIRNPTRVIIEEIDDDENDIDAVCSQESSK TSDSSITSADOSKTMLLDPFNTVLMNEOVDSOMVKGKGHIPYTDDLNDLSQGISMVSSAS THCELNLNEVPPEVELCSHOODPESTIOTODOOESTRTEDVKKNRKKPTTSKPKKKSKES AKSTOKKSVDWDSLRKEAESGGRKRERTERTMDTVDWDALRCTDVHKIANIIIKRGMNNM LAERIKAFLNRLVKKHGSIDLEWLRDVPPDKAKEYLLSINGLGLKSVECVRLLSLHQIAF PVDTNVGRIAVRLGWVPLOPLPDELOMHLLELYPVLESVOKYLWPRLCKLDOKTLYELHY HMITFGKVFCTKVKPNCNACPMKAECRHYSSARASARLALPEPEESDRTSVMIHERRSKR KPVVVNFRPSLFLYOEKEOEAORSONCEPIIEEPASPEPEYIEHDIEDYPRDKNNVGTSE DPWENKDVIPTIILNKEAGTSHDLVVNKEAGTSHDLVVLSTYAAAIPRRKLKIKEKLRTE HHVFELPDHHSILEGFERREAEDIVPYLLAIWTPGETVNSIQPPKQRCALFESNNTLCNE NKCFOCNKTREEESOTVRGTILIPCRTAMRGGFPLNGTYFQTNEVFADHDSSINPIDVPT ELIWDLKRRVAYLGSSVSSICKGLSVEAIKYNFOEGYVCVRGFDRENRKPKSLVKRLHCS HVAIRTKEKTEE

SEO ID NO:12

>DMT3(1DMT3) novel 375 amino acid amino terminus;

MEVEGEVREKEARVKGRQPETEVLHGLPQEQSIFNNMQHNHQPDSDRRRLSLENLPGLYN
MSCTQLLALANATVATGSSIGASSSSLSSQHPTDSWINSWKMDSNPWTLSKMQKQQYDVS
TPQKFLCDLNLTPEELVSTSTQRTEPESPQITLKTPGKSLSETDHEPHDRIKKSVLGTGS

PAAVKKRKIARNDEKSQLETPTLKRKKIRPKVVREGKTKKASSKAGIKKSSIAATATKTS EESNYVRPKRLTRRSIRFDFDLQEEDEEFCGIDFTSAGHVEGSSGEENLTDTTLGMFGHV PKGRRGQRRSNGFKKTDNDCLSSMLSLVNTGPGSFMESEEDRPSDSQISLGRQRSIMATR PRNFRSLKKLLQRII

SEQ ID NO:13

5

	>DMT3	(1DMT3) nu	cleotide s	sequence f	rom BAC T2	2K18 (gi 1	2408726);
	53341	aatcaagtac	taatgcagat	ttaagggggg	tgtattgacg	gcgttaaaac	ggtttctcaa
	53401	cggaatcgta	cgtagtcaca	cgtgatttta	ttgtttaccc	cggattggtc	atgcgttcct
10	53461	tcttttccac	ttgcgcggac	cactcaatga	cactctcttc	ttttgtagca	gtggcccgac
	53521	accagaatgc	agcatttaat	ctctcaaatt	accattttgc	tcctacctct	tttacccctt
	53581	ttggtatttt	gtgtctttt	tctttctatt	tcgtgtgaaa	aaggatctct	tccttaatcg
	53641	tattatttct	tccgatatct	acttttattc	tgttttctat	ttttggtagg	ttacatcttt
	53701	tttataaaga	aaatatgagc	taacacgaca	ttagtgttgt	taaccaaaga	attggaaaaa
15	53761	agttataaga	gagataataa	gattctctta	cagagactca	cttcagtgaa	aaaggaagaa
	53821	gcaagtggtt	cccttaaggg	aaaaaaagt	cacgtacgtt	catatacaac	tttaatacgt
	53881	actgtgtaac	tcaatagatc	gtgcagtaat	attcagtcgt	attagtaaga	aggaatttat
	53941	ttgctaagta	aactcaagcc	tcctttttct	ctttttttc	tttttagtaa	aaattaggct
	54001	agtgttttt	ttgactcagc	aacactctgc	ttaaatttag	gagtaatttg	acctattcct
20	54061	acgagtttct	aagtgaattc	tgttggggtc	aaagaagcaa	ctagttgaat	tagtggaaaa
	54121	tegttteett	tctttacgca	tagttcacgt	tggacactca	gtctcaatgc	tttcacgttt
	54181	cacgtagcaa	caacatatat	tcatcagttt	gtgatcgtgc	catcgtggat	aagttgcaat
	54241	tcagtgaaac	tctgcaccac	tttgtgcaat	tatttggccg	tctaatctat	ttgtgagaat
. 0.5	54301	tttacaatct	aattgttcta	ttatttcatt	tacttgtcat	caatttatta	tatttgtagc
25	54361	caatgaacgt	tgtaattaaa	gaaccaaaat	aaattaatat	cttgaaattt	gtaacagtca
;	54421	ctagaagctg	atttcttatt	aattgtatca	ctaaagtatt	attaaaaacg	gttacaaatt
_	54481	atgataatta	tatatttaat	aaatttcgtg	tgtcacattt	cttttaaact	acaattatga
	54541	atatctaaaa	ctcattcatg	-catatcttaa	aatttgaatt	caaaactttc	ttatcttatc
20	54601	tttaggttct	taattaacag	tcactaaaaa	tagtcaaagt	tttgaagttt	atgaaaaaag
30	54661	ataagagtat	aattaatgga	tacgcctcgt	aacaaattct	tgtaaagtat	agataatata
	54721	catttgttaa	atatgacacg	tgtttatttt	ttttttaaat	atgatcaaaa	tatattttaa
	54781	ctacctagat	ggtatgtatg	tctccaattt	tgaataacaa	gtcaattgtt	attagaaatg
						tgaaaaacaa	
2.5	54901	ttttggagaa	aattaaggac	tggatttgca	aaaacgaaaa	agtaacttca	tgtatattgt
35	54961	cttccttata	gtctctataa	actattatct	caaattttgt	ctggactctg	aaactcacaa
	55021	gacttgactc	tggcttactt	ggcttcatct	ttttctctct	ggtaatctct	cctgcaactt
	55081	caagctttca	ttttcaaata	aatgtaatca	aatctgttat	tttcactcaa	gaactaattg
	55141	agttctctat	ccctttcaat	tgaaattgac	attaaaatga	aaagattttg	aggaggtttc
40	55201	acctaccaca	accgaatcac	ttetttetee	aaatattgtt	tctttcagtg	gccaagaatc
40			_			attgaacaga	
						tgagagagaa	
						tgccacaaga	
					_	ggttttgtga	
15		/-		_		gattctgttt	_
45						caaaacatga	
			_	_		tggttttggc	
						taaggagaga	
	55741	aactgttatt	gatcatacag	gaggaggctt	agtcttgaaa	acttacctgg	actatacaac

55801 atgtettgta cacaactett ggetetggee aatgeeaeag tegecaeagg tteateaatt

	55861	ggtgcatcat	catcatcgtt	aagctctcag	catccaacgg	attcttggat	taatagctgg
	55921	aagatggact	ctaatccgtg	gactttgagt	aaaatgcaaa	aacaacaatg	tgagtaaaat
	55981	ttgttcctga	atttgtagga	tcttttaaga	gaaagtaagc	gtttatgtgt	agattaagtc
•	56041	agactgaaat	cgattatctc	ataataagtt	ctcagtgatc	tctcaaatca	tgaattttat
5	56101	gtttacctga	tatcaacttc	ttgtcttggt	gaaccacaga	tgatgtgtca	actccgcaga
	56161	agtttctttg	tgaccttaat	cttacacctg	aagagttggt	gagcaccagt	acgcaacgaa
	56221	cagaacctga	gtctcctcaa	ataactttaa	agacaccagg	aaaaagtctg	tctgaaactg
	56281	atcatgagcc	tcacgaccgt	atcaagaagt	ctgttcttgg	aactggatct	cctgcagcag
	56341	taaagaaaag	aaagatagca	agaaatgatg	agaaatctca	gctggaaaca	ccaacactaa
10	56401	agagaaaaaa	gatcaggcca	aaggttgtcc	gtgaaggcaa	aacaaaaaa	gcatcatcta
	56461	aagcagggat	taaaaaatcc	tctattgctg	ctactgctac	taaaacttct	gaagagagca
	56521	attatgttcg	gccaaaaaga	ttaacgagaa	gatctatacg	attcgacttt	gaccttcaag
	56581	aagaagatga	ggaattttgt	ggaatcgatt	tcacatcagc	aggtcacgta	gagggttctt
	56641	caggtgaaga	aaatctaacc	gatacaacac	tgggaatgtt	tggtcacgtc	ccaaagggaa
15	56701	gaagagggca	aagaagatcc	aatggcttta	aaaaaaccga	caatgattgc	ctcagttcta
	56761	tgttgtctct	tgtcaatacc	ggaccaggaa	gtttcatgga	atcagaagaa	gatcgtccga
	56821	gtgattcaca	aatttctctg	ggaagacaga	gatccattat	ggcaaccaga	ccgcgtaact
	56881	tccgatcgtt	aaagaaactt	ttacaaagga	ttataccaag	caaacgtgat	agaaaaggat
	56941	gtaagcttcc	tcgtggactt	ccgaagctta	ccgtcgcatc	caagttgcaa	ctaaaagtgt
20	57001	ttagaaagaa	gcggagtcaa	agaaaccgtg	tagcaagcca	gttcaatgca	aggatattgg
•	57061	acttgcagtg	gcgacgccaa	aatccaacag	gtgataaaca	cacaagcaac	tttcatctat
	57121	aatattttc	ttagatttct	atcttttgaa	ttaatactag	ttttacaaaa	tgcaggtaca
	57181	tcgctagctg	atatatggga	aagaagtttg	actattgatg	ctatcactaa	gttgtttgaa
	57241	gaattagaca	tcaacaaaga	gggtctttgc	cttccacata	atagagaaac	tgcacttatt
25	57301	ctatacaaaa	agtcgtatga	agagcaaaag	gcaatagtga	agtatagcaa	gaagcagaaa
	57361	ccgaaagtac	aattggatcc	tgaaacgagt	cgagtgtgga	aactcttaat	gtcaagtatc
	57421	gactgtgacg	gtgttgatgg	atcagatgag	gaaaaacgta	aatggtggga	agaggagagg
	57481	aacatgttcc	atggacgtgc	aaactcgttc	attgcgcgaa	tgcgtgttgt	ccaaggtatt
	57541	atttattgct	ttagttatga	cattgttgtg	tggctttata	ccttagatct	ttctttctt
30	57601	cttttttgta	tccaaagcaa	catggtctta	aatcaagctt	atcactgcag	gcaatagaac
	57661	tttctcacct	tggaaagggt	cagtagtgga	ttcagtagtg	ggagttttcc	taacccagaa
	57721	tgtcgcagac	cattcatcaa	ggtatatgca	ttcaagagat	ttctaataag	tagaagatat
	57781	atgcaacaga	gtggtttaga	aattataact	tgttcacttt	tgcagttctg	catatatgga
	57841	tttagctgct	gagtttcctg	tcgagtggaa	cttcaacaag	ggatcatgtc	atgaagagtg
35	57901	gggaagttca	gtaactcaag	aaacaatact	gaatttggat	ccaagaactg	gagtttcaac
			cgcaatccaa				
			gtttgtagtc				
			aaaacgatgc		_		
			atggtaaaag				
40		_	atttcgatgg				_
			gaagtagagt				
		•	caagagagca				
			ccaaagaaaa				
			agtttgagaa				
45			gacacagttg				
		-	atcaaacgag			_	
			ctatatatac				
			caggccttct				
			gatgttcctc				
50			gcctctggtt		-		
-			atgagagata				
						~ CCC	J

	58921	agcataaacg	gattaggatt	gaagagtgtg	gagtgtgtta	gacttttgtc	actacatcag
	58981	attgcattcc	ctgtaagtca	atgaaggata	ctgaatactc	agaccctaat	gaatgtggaa
	59041	cagatacatt	aatagttacg	tatttttaca	aatgcaggtt	gacacgaatg	tcggacgcat
	59101	agctgtaaga	ctaggatggg	ttcccttaca	gccattgccc	gacgagctgc	aaatgcatct
5	59161	tttagagttg	taagaaaaaa	aaattaaaga	tcattcttca	atcatgaaag	ggaacatgag
	59221	aaatttacag	tagttccctt	taattctatt	caggtaccca	gttctagagt	cagttcaaaa
	59281	gtacctctgg	ccacgcctct	gcaagcttga	ccaaaaaacc	ttgtaagtaa	attacattag
	59341	catcaaccat	tactctagac	ccttaaactt	ctctaactaa	ctctaactgt	atcatacaat
	59401	tctaggtacg	agctgcatta	ccacatgata	acatttggaa	aggtacctca	aacaaatttc
10	59461	aagtgtttgt	ggaatgaaaa	catcttaaag	tggcttttcc	tattttgcag	gtcttttgca
	59521	caaaagtaaa	acccaattgc	aatgcatgtc	caatgaaggc	ggagtgtcga	cattactcta
	59581	gtgcacgtgc	aaggttaaac	cccacaaaat	tctttgttat	tgccattaac	atgaaaaaaa
	59641	aaacactagc	ttaaagagaa	agagatctgc	tcaaaatagt	cattttaatg	gttgtatgtt
	59701	ctaaatgctt	gtgttatatc	gcagcgcacg	gcttgcttta	ccagaaccag	aggagagtga
15	59761	cagaacaagt	gtaatgatcc	atgagaggag	atctaaacgc	aagcctgttg	tggttaattt
	59821	tcgaccatcc	ttatttcttt	atcaagaaaa	agagcaagaa	gcacaaagat	cccaaaactg
	59881	tgaaccaatc	attgaggaac	cagcatcacc	agaaccagag	tatatagaac	atgatattga
	59941	agactatcct	cgggacaaaa	acaacgttgg	aacatcagag	gatccttggg	aaaataagga
	60001	cgtaattcct	accatcatcc	tcaacaagga	agctggtaca	tcacatgatt	tggtggtcaa
20	60061	caaggaagct	ggtacgtcac	atgatttggt	ggtactaagc	acatatgcag	cagcaatacc
	60121	tagacgtaaa	ctcaagatca	aggaaaagct	acgcacagag	caccacgtgt	gagttgccac
	60181	tttcaatttt	ttcttctatt	ataccctaaa	ccgtaaaatt	tgagactttc	ctcagcattt
	60241	atctcatact	aattctcttt	tacagatttg	agctccctga	tcaccattcc	attctagaag
	60301	gggttagtaa	ctcttgcaaa	atgatttagc	aagaatttt	ctacttattc	ccgccttaaa
25	60361	aactgtttga	ttatctttt	ttacagtttg	agaggcgaga	agctgaggat	atagtccctt
	60421	acttgttagc	catttggacg	ccaggtaaga	agaaataggc	acacaataaa	atctgattat
	60481	gatttttctt	ttcaagaata	ccgctatatt	tttacgagtt	ttcatcctta	gatgtatatg
	60541	actaatgtct	aacaagtgat	tgtaatattt	ttccatacca	ggtgaaaccg	tgaattccat
	60601	tcaaccgcca	aaacaaagat	gtgctttatt	tgaaagcaat	aatacattat	gcaacgaaaa
30	60661	caaatgtttt	caatgcaaca	agacacggga	agaggaatca	cagactgtac	gaggaactat
	60721	attggtaaga	ttctggtgga	caattttcaa	gagaatatct	ctaagtagaa	atataaggaa
	60781	ggtataaaaa	tgactaattt	gtttgttaac	agataccttg	cagaacagca	atgagaggtg
	60841	gattcccttt	gaatggcaca	tacttccaaa	ctaatgaggt	aattttccca	aaaatgaatt
	60901	taacttaaac	aaatgatcaa	aagcaacatt	ctcgtcaaag	ctcgatttgg	actatacttg
35	60961	tgcaggtttt	tgctgaccat	gactctagca	taaaccctat	cgacgtccca	acagaactga
	61021	tatgggatct	aaaaagaaga	gtcgcatact	taggatcctc	tgtatcctcg	atttgtaaag
	61081	gtaaattttc	aaaacaaaac	tgtcgattta	tgcatgtgtt	tggatatata	aatccaaggt
	61141	cttgtctcaa	tatgtttttc	tcatttttt	aggtttatca	gtggaagcca	taaaatacaa
	61201	tttccaggaa	ggtatgctaa	tatgtcttac	actgaaaaca	cctttagtat	caaacattga
40	61261	attcatgaaa	agaacaaaca	atagtatcaa	aatcagtcac	gatgtttttg	ctttggcgat
	61321	gtaagatgtt	gataggaaag	tatagaagat	atagcttaag	ttggttaata	ctgtttttat
	61381	agagctttga	ggtggggttt	gactagcatt	gtaatatata	tgcaggatat	gtctgtgtaa
	61441	ggggattcga	cagggagaat	cgtaagccaa	agagtctagt	gaaaagactg	cattgttctc
4.5	61501	acgtagcaat	cagaactaaa	gagaagacag	aggaatgaaa	ccttccagat	tgcattaaca
45	61561	tgttagacat	atttgattca	ttggtttagg	gtttacatca	ccaaggtcat	agaggatett
	61621	agcttttcat	taacttttaa	attcatgcaa	ctctttttag	gtgtttcttt	ttgttccttg
	61681	ccatagtttt	gggcaatgga	tggatgttct	ttgcaaactc	aggtttttg	tagtcattaa
	61741	cagaaatttg	cagcactaat	tcatctttcc	tattatctat	caaagctctc	agtgtttctc
50	61801	cataacttga	tgagatttag	tcactctcaa	gctaattcag	tctggtccta	atttcaatca
50	61861	gatttggtaa	aggaacaact	gcaattgcta	agtacaaatc	gatccagatt	tcaaacaagt
	61921	tccaggttta	atccaaatca	tcacattcaa	tcaaagacca	aactagaatt	caaaacatat

15

20

25

61981 aatototgat toagattoaa gaaagacaaa goatgagaca toattotgoa agttaaccaa 62041 ttooggttat totogaatoo taotgaatta agoatoaato atotaaagga acttoataag

5 SEQ ID NO:14

Arabidopsis thaliana DMT4

>DMT4(1DMT4);

MEFSIDRDKNLLMVVPETRIKTKQFEKVYVRRKSIKLPQNSMVHNTLIKMARQRIQKSMK ESVMNQHIFKNFDSYLSVIYHPCCFVINNSQTTHKKKEKKNSKEKHGIKHSESEHLQDDI SORVTGKGRRRNSKGTPKKLRFNRPRILEDGKKPRNPATTRLRTISNKRRKKDIDSEDEV I PELATPTKESFPKRRKNEKI KRSVARTLNFKQEI VLSCLEFDKI CGPI FPRGKKRTTTR RRYDFLCFLLPMPVWKKOSRRSKRRKNMVRWARIASSSKLLEETLPLIVSHPTINGQADA SLHIDDTLVRHVVSKQTKKSANNVIEHLNRQITYQKDHGLSSLADVPLHIEDTLIKSASS VLSERPIKKTKDIAKLIKDMGRLKINKKVTTMIKADKKLVTAKVNLDPETIKEWDVLMVN DSPSRSYDDKETEAKWKKEREIFOTRIDLFINRMHRLOGNRKFKOWKGSVVDSVVGVFLT ONTTDYLSSNAFMSVAAKFPVDAREGLSYYIEEPODAKSSECIILSDESISKVEDHENTA KRKNEKTGI I EDE I VDWNNLRRMYTKEGSRPEMHMDSVNWSDVRLSGQNVLETTI KKRGQ FRILSERILKFLNDEVNONGNIDLEWLRNAPSHLVKRYLLEIEGIGLKSAECVRLLGLKH HAFPVDTNVGRIAVRLGLVPLEPLPNGVQMHQLFEYPSMDSIQKYLWPRLCKLPQETLYE LHYOMITFGKVFCTKTIPNCNACPMKSECKYFASAYVSSKVLLESPEEKMHEPNTFMNAH SQDVAVDMTSNINLVEECVSSGCSDQAICYKPLVEFPSSPRAEIPESTDIEDVPFMNLYQ SYASVPKIDFDLDALKKSVEDALVISGRMSSSDEEISKALVIPTPENACIPIKPPRKMKY YNRLRTEHVVYVLPDNHELLHDFERRKLDDPSPYLLAIWQPGETSSSFVPPKKKCSSDGS KLCKIKNCSYCWTIREONSNIFRGTILVFADHETSLNPIVFRRELCKGLEKRALYCGSTV TSIFKLLDTRRIELCFWTGFLCLRAFDRKQRDPKELVRRLHTPPDERGPNGFHIVVVDEK EESPRVGLMVMPGFWIGGSVIONRVYVSGVKVLE

SEQ ID NO:15

>DMT4 novel 372 amino acid NH2 terminus;


30 MEFSIDRDKNLLMVVPETRIKTKQFEKVYVRRKSIKLPQNSMVHNTLIKMARQRIQKSMK
ESVMNQHIFKNFDSYLSVIYHPCCFVINNSQTTHKKKEKKNSKEKHGIKHSESEHLQDDI
SQRVTGKGRRRNSKGTPKKLRFNRPRILEDGKKPRNPATTRLRTISNKRRKKDIDSEDEV
IPELATPTKESFPKRRKNEKIKRSVARTLNFKQEIVLSCLEFDKICGPIFPRGKKRTTTR
RRYDFLCFLLPMPVWKKQSRRSKRRKNMVRWARIASSSKLLEETLPLIVSHPTINGOADA

SLHIDDTLVRHVVSKQTKKSANNVIEHLNRQITYQKDHGLSSLADVPLHIEDTLIKSASS VLSERPIKKTKD

SEQ ID NO:16

SEQ ID NO.10								
5	>DMT4	nucleotide	sequence B	AC F28A23 (gi 7228244);	;		
	14881	gctatggatg	tcaacagaga	gaattacgaa	ttgggtttac	cgatcattga	gaaagccggc	
	14941	gttgctcaca	agatcgactt	cagggaaggc	cctgctcttc	ccgttcttga	tgaaatcgtt	
	15001	gctgacgtaa	gcattcttct	ttctgacgta	attaacaaaa	aagatgatga	agataatgaa	
	15061	ataattaaaa	actcatggcc	taattaggtt	gatttaatat	cttgatgaga	atttctgtat	
10	15121	acgcaaattt	gtttcctttt	tcatagaaga	aagtgtggta	actgattatt	gtgtgtggtt	
	15181	gggtgcagga	gaagaaccat	ggaacatatg	actttatatt	cgttgatgct	gacaaagaca	
	15241	actacatcaa	ctaccacaag	cgtttgatcg	atcttgtgaa	aattggagga	gtgattggct	
	15301	acgacaacac	tctgtggaat	ggttctgtcg	tggctcctcc	tgatgcacca	atgaggaagt	
	15361	acgttcgtta	ctacagagac	tttgttcttg	agcttaacaa	ggctcttgct	gctgaccctc	
15	15421	ggatcgagat	ctgtatgctc	cctgttggtg	atggaatcac	tatctgccgt	cggatcagtt	
	15481	gatttgactc	ctccctactc	tgagtttgtc	cacagtggat	tactttccat	cttcttatac	
	15541	ctttcaatcg	cattttcacc	aaccactaaa	atggaccttt	ttatgtattt	gtgttaagta	
	15601	atatctccat	tgtccttgtt	ttgctttctt	ctgaacaaag	aaataatatg	taccttactt	
	15661	ttcttcttgg	tctcgttctt	ttgtttttct	ccatgataca	acatctaaag	aaattatttg	
20	15721	tgtcacagca	acgtaagtcg	ataaaattag	ttgaacatat	tgagaaaaag	ttatcataga	
	15781	ccttcaattg	ttgaaagtcg	atgttggtat	ttgtcaattg	atattagatt	accaaataaa	
	15841	tattagacag	taagaaacga	acaaagtagg	aagatgtagg	tcaccggtct	ttgaaaattt	
	15901	atcagataga	attcataata	cacagttagg	tagtttcagt	tgagagttaa	aagggaaaaa	
25	15961	tatgtaattg	tgtgtgataa	atacgtcaaa	aattagttga	tgagcaaaat	cgtaaacaaa	
25	16021	aatactttt	tgcattagtt	ttgttggatt	ccctataaat	acgggttccc	atatctaact	
	16081	cgtagttagc	ataattataa	gcaacaaata	aacacaaaat	actgaattta	gaaattttcc	
	16141	agaaaattaa	ttagagattt	tacattattt	ttacaaactt	tagtgaatta	tttcttaaac	
-	16201	gtatgttagt	tatttattaa	ctgaagtttc	acatatttga	tagaataaca	tttaaataaa	
20	. 16261	aaaatttgaa	gtaaggttag	aatgttctta	taatacttta	taacttttt.	aaaaggtaca	
30	16321	agccaaaatt	atcgcaaatg	taaataataa	atcattgtaa	aaatcttaaa	ctaattaaaa	
	16381	gatctaacgc	aatctaaaca	aagatttggt	atcatcgccc	atttatgttt	tgatataatc	
							ttcttgttaa	-
	16501	tgtaatcaac	tcaccattat	tttaattatt	taaaatatgg	gttaatatct	cttaatcata	
25				catttatgaa				
35				taatattaaa		_		
				ctttttcca				
		_		tttcttcttt			=	
			_	aaaatatttt	_		_	
40				cattatacta				
40				gacaaaaatc				
			_	gtttatgtga		=		
			_	atcaaagatc				
				aatctctaat				
45		Ā		atgagagtat		_		
43				gactaattat	_		_	
				attttattga			_	
		_	Ī	tatacttttg				
				gttcattgat			_	
50				atttgtttcg				
50	17521	agatggttca	caacacactt	atcaaaatgg	cgagacaaag	aattcaaaag	agcatgaaag	

	17581	aaagtgtgat	gaatcagcac	atcttcaagg	taaataattt	taaattcatt	cttaaaaaag
	17641	ttagcttatt	ggtaagttca	ttacaattta	tatttaacca	tcgtcacttt	ttatttaacg
	17701	agtttgataa	gcattttcaa	aacctgtcct	tcatctgccg	atgcagatgt	ggttatgttc
	17761	atctttgatt	ttattgattg	aggattttt	cagaatttcg	attcatactt	gtctgtaata
5	17821	tatcatccat	gttgttttgt	aatcagttaa	ttcacttatt	ttattttaa	cttttattgt
	17881	aacagataat	tcacaaacca	cccataaaaa	aaaggagaag	aagaattcaa	aagaaaagca
	17941	tggaataaag	cattctgaat	cagaacatct	tcaaggtaaa	tacttttgaa	ttcattcatt
	18001	aaaaaacag	tttatttgta	agttcattac	agtttatata	tatttaaatt	gtttatgata
	18061	atgtatttt	gcacaatcga	ctaatcatta	cccactcatt	catttatatt	ttattttatg
10	18121	gtgaaagatg	atatttcgca	acgtgttacc	ggaaaaggaa	ggagaaggaa	ttcaaaaggg
	18181	acaccaaaaa	aactgaggtt	taataggcct	cggatcttgg	aagacggaaa	gaaaccaaga
	18241	aatcccgcca	ccactcgact	gagaactata	tccaacaaga	ggaggaaaaa	ggacatagac
	18301	agtgaagatg	aagttatacc	agagcttgca	actccaacaa	aggaaagctt	tccaaagaga
	18361	agaaagaacg	agaagattaa	gagatccgtg	gctcggactt	taaattttaa	gcaagaaatt
15	18421	gttctgagtt	gtcttgagtt	cgačaagatt	tgtggaccaa	tttttccaag	agggaaaaag
	18481	aggaccacca	cacgacgcag	atatgatttc	ctttgttttt	tacttccgat	gcctgtttgg
	18541	aaaaaacaat	caagaaggtc	taagcgtagg	aaaaatatgg	tcagatgggc	tagaattgct
	18601	tcttcttcaa	aactgctaga	agaaactttg	cctttaatag	taagtcatcc	gactattaat
	18661	ggacaagcag	atgcttcttt	acacattgat	ggtaatcgag	tttttttt	gttaatttat
20	18721	ctgttacatc	aaaattgttt	atgcttatat	ctaaagtatc	attgtgtatt	attttttgca
	18781	gacacactcg	tgagacatgt	agtctcaaag	caaaccaaga	aaagtgctaa	caatgtcatt
	18841	gagcatttaa	atcgacaaat	aacttatcag	aaagatcacg	gtctctcatc	tctggcagat
·	18901	gttcctttgc	acattgaagg	taatctagtc	ttatttttgt	tctttttaa	tatattgatt
-	18961	aaaaagattg	tgatatattt	atttaatata	tttttgttat	attatatcta	tattttattg
25	19021	tttgtacttt	ttttttgtag	atacactaat	aaaatcggct	agttctgtac	tttcagaacg
	19081	acccatcaag	aaaactaagg	atattgctaa	gttaatcaaa	gatatgggaa	gattaaagat
	19141	caataaaaag	gtaacaacga	tgatcaaagc	tgacaagaaa	ctcgttacgg	caaaggttaa
•			gagaccatta				
•	19261	atcatatgac	gataaggaga	cggaggccaa	atggaaaaaa	gaaagagaga	tttttcaaac
30	19321	ccggatagat	cttttcatta	accggatgca	tcgcttacaa	ggtacattat	tgttattatc
	19381	attattgtta	ttatgatcta	tttatacttg	tattctaaat	tagcttacat	atatatataa
	19441	ggaatccaag	tataagtgag	tatgctaagt	atatgatcat	tttttgaaat	tatgtttcct
			ttaaatgatt				
			tcagtggttg				
35			tgtttaaatt				
			tgagtttcta				
			aaggtctatc				
			tatctgatga				
	19861	aggaaaaacg	agaaaaccgg	tattatagaa	gatgagatag	ttgactggaa	caatcttaga
40	19921	aggatgtaca	cgaaagaagg	atctcgtccc	gaaatgcata	tggactctgt	taattggagt
			tatctggcca				
	20041	aggattcttt	cagaaagaat	attggtaaga	aaaacaaac	ttctaatgaa	ctttgtgaat
	20101	aatttattca	aatgatttaa	gactaacact	tttttttt	tecttgtttt	ctcaagaaat
			tgaagttaac	-			_
45		_	agtgaagtat		_		
			ataaactctt		-	-	
			aaaagtgctg				
			tattattatg				
			taatgtgtag				-
50			tgaaccttta			_	
			tttgattaag		_		
]				



5

10

15

20

25

30

RICE(Oryza sativa) DMT1

>DMTRICE(1DMTRICE);

MQDFGQWLPQSQTTADLYFSSIPIPSQFDTSIETQTRTSAVVSSEKES ANSFVPHNGTGLVERISNDAGLTEVVGSSAGPTECIDLNKTPARKPKKKKHRPKV LKDDKPSKTPKSATPIPSTEKVEKPSGKRKYVRKKTSPGQPPAEQAASSHCRSELK SVKRSLDFGGEVLQESTQSGSQVPVAEICTGPKRQSIPSTIQRDSQSQLACHVVSST SSIHTSASQMVNAHLFPPDNMPNGVLLDLNNSTSQLQNEHAKFVDSPARLFGSRI RQTSGKNSLLEIYAGMSDRNVPDLNSSISQTHSMSTDFAQYLLSSSQASVRETQM ANOMLNGHRMPENPITPSHCIERAALKEHLNHVPHAKAAVMNGQMPHSYRLAQ NPILPPNHIEGYQVMENLSELVTTNDYLTASPFSQTGAANRQHNIGDSMHIHALD PRRESNASSGSWISLGVNFNQQNNGWASAGAADAASSHAPYFSEPHKRMRTAYL NNYPNGVVGHFSTSSTDLSNNENENVASAINSNVFTLADAQRLIAREKSRASQRM ISFRSSKNDMVNRSEMVHQHGRPAPHGSACRESIEVPDKQFGLMTEELTQLPSMP NNPQREKYIPQTGSCQLQSLEHDMVKGHNLAGELHKQVTSPQVVIQSNFCVTPP DVLGRRTSGEHLRTLIAPTHASTCKDTLKALSCQLESSRDIIRPPVNPIGPSSADVP RTDNHQVKVSEETVTAKLPEKRKVGRPRKELKPGEKPKPRGRPRKGKVVGGELA SKDSHTNPLQNESTSCSYGPYAGEASVGRAVKANRVGENISGAMVSLLDSLDIVI QKIKVLDINKSEDPVTAEPHGALVPYNGEFGPIVPFEGKVKRKRSRAKVDLDPVT ALMWKLLMGPDMSDCAEGMDKDKEKWLNEERKIFQGRVDSFIARMHLVQGDR RFSPWKGSVVDSVVGVFLTQNVSDHLSSSAFMALAAKFPVKPEASEKPANVMFH TISENGDCSGLFGNSVKLQGEILVQEASNTAASFITTEDKEGSNSVELLGSSFGDG VDGAAGVYSNIYENLPARLHATRRPVVQTGNAVEAEDGSLEGVVSSENSTISSQN SSDYLFHMSDHMFSSMLLNFTAEDIGSRNMPKATRTTYTELLRMQELKNKSNETI ESSEYHGVPVSCSNNIQVLNGIQNIGSKHQPLHSSISYHQTGQVHLPDIVHASDLE QSVYTGLNRVLDSNVTQTSYYPSPHPGIACNNETQKADSLSNMLYGIDRSDKTTS LSEPTPRIDNCFQPLSSEKMSFAREQSSSENYLSRNEAEAAFVKQHGTSNVQGDN TVRTEQNGGENSQSGYSQQDDNVGFQTATTSNLYSSNLCQNQKANSEVLHGVSS NLIENSKDDKKTSPKVPVDGSKAKRPRVGAGKKKTYDWDMLRKEVLYSHGNKE RSONAKDSIDWETIROAEVKEISDTIRERGMNNMLAERIKDFLNRLVRDHGSIDLE WLRYVDSDKAKDYLLSIRGLGLKSVECVRLLTLHHMAFPVDTNVGRICVRLGW VPLQPLPESLQLHLLEMYPMLENIQKYLWPRLCKLDQRTLYELHYQMITFGKVFC TKSKPNCNACPMRAECKHFASAFASARLALPGPEEKSLVTSGTPIAAETFHQTYIS SRPVVSQLEWNSNTCHHGMNNRQPIIEEPASPEPEHETEEMKECAIEDSFVDDPEE

IPTIKLNFEEFTQNLKSYMQANNIEIEDADMSKALVAITPEVASIPTPKLKNVSRLR TEHQVYELPDSHPLLEGFNQREPDDPCPYLLSIWTPGETAQSTDAPKSVCNSQEN GELCASNTCFSCNSIREAQAQKVRGTLLIPCRTAMRGSFPLNGTYFQVNEVFADH DSSRNPIDVPRSWIWNLPRRTVYFGTSIPTIFKGLTTEEIQHCFWRGFVCVRGFDRT SRAPRPLYARLHFPASKITRNKKSAGSAPGRDDE

SEQ ID NO:18

5

>DMTRICE novel 723 amino acid NH2 terminus;

MQDFGQWLPQSQTTADLYFSSIPIPSQFDTSIETQTRTSAVVSSEKESANSFVPHNGTGLVERISNDAGLTEVVGSSAGP
TECIDLNKTPARKPKKKKHRPKVLKDDKPSKTPKSATPIPSTEKVEKPSGKRKYVRKKTSPGQPPAEQAASSHCRSELKS
VKRSLDFGGEVLQESTQSGSQVPVAEICTGPKRQSIPSTIQRDSQSQLACHVVSSTSSIHTSASQMVNAHLFPPDNMPNG
VLLDLNNSTSQLQNEHAKFVDSPARLFGSRIRQTSGKNSLLEIYAGMSDRNVPDLNSSISQTHSMSTDFAQYLLSSSQAS
VRETQMANQMLNGHRMPENPITPSHCIERAALKEHLNHVPHAKAAVMNGQMPHSYRLAQNPILPPNHIEGYQVMENLSEL
VTTNDYLTASPFSQTGAANRQHNIGDSMHIHALDPRRESNASSGSWISLGVNFNQQNNGWASAGAADAASSHAPYFSEPH
KRMRTAYLNNYPNGVVGHFSTSSTDLSNNENENVASAINSNVFTLADAQRLIAREKSRASQRMISFRSSKNDMVNRSEMV
HQHGRPAPHGSACRESIEVPDKQFGLMTEELTQLPSMPNNPQREKYIPQTGSCQLQSLEHDMVKGHNLAGELHKQVTSPQ
VVIQSNFCVTPPDVLGRRTSGEHLRTLIAPTHASTCKDTLKALSCQLESSRDIIRPPVNPIGPSSADVPRTDNHQVKVSE
ETV

20 SEQ ID NO:19

>DMTRICE nucleotide sequence from PAC PO489G09;

```
10261 aaatattgct taaatggata taaagttgaa aaatgtactt gagggaagtt gtaggtgcac
          10321 gtggggtccc acaatttttc ttcactagtg cacctttagt tatatatttt ttgcgcaaga
          10381 ggacaaaggc gctccgtgta attttgagta agggccggcg ggatatttat ttgtgtaaag
25
          10441 gacctagcca agaaaagcat gatagtgcat atgtateett tettttett ttettttgtt
          10501 ttcataactg tcttacagaa tttcatgttg gctggtgaca cttgtctcac tcattatttg
          10561 gtatattttg actaaatgca acgtgttggt gctcggtagt ttatatttgt ttttacgcat
          10621 tetteattga etgtatgtat ttgatgttga taccetggge tgtettattt tataggtgga
          10681 tgctgggagg ccacatagga ggcctgtgtg atccaagtgt gctgctcctg agttgaaatt
30
          10741 gcatagccat atagcaacta ctggtgtaaa cttgagagat gaagtagtga aaggaaatat
          10801 gcaggatttt ggacaatggc tgcctcaatc tcagaccact gccgatctat atttctccag
          10861 tattccaata ccatcacagt tcgatacttc catagagacg cagactagaa cttctgcagt
          10921 tgtatcgtca gagaaagaat ctgctaattc gttcgtccct cataatggta ctgggcttgt
          10981 tgaacgcatt agcaatgatg ctgggctaac tgaagtagtt ggaagtagtg ctggaccaac
35
          11041 tgaatgtatt gacttgaaca agacaccagc acggaaaccc aagaagaaaa agcacaggcc
          11101 aaaggtgcta aaggacgata aaccatcgaa gacacctaaa tctgctactc caataccttc
          11161 aacagaaaag gtagaaaaac catctggaaa gagaaaatat gtccgcaaga agacatctcc
          11221 aggccaacct cctgcagaac aggcagctag ctcacactgc agatctgagc tgaagtcagt
          11281 taaacgaagt ttggactttg gtggagaagt actgcaagag agtacacaat ctggatctca
40
          11341 agttccggtg gcagaaatat gtactggtcc caagcgtcaa tcaatacctt ctaccatcca
          11401 aagagatteg caaagecagt tggettgeca egtggtttet ageaceaget caatteacae
          11461 ttcagctagt cagatggtta atgcacattt gtttcctcct gataacatgc caaatggagt
          11521 attgcttgac ctcaataatt ctactagtca gttacaaaac gaacatgcta aatttgtgga
          11581 cagtccggca cgtctttttg gttccagaat aagacagaca tcaggtaaaa attctttgct
45
          11641 agaaatctat gctggcatgt cagatagaaa tgtacctgat ctcaacagtt caatcagtca
          11701 gacgcatage atgtctactg attttgctca atacttgctt tcatcctcac aagcttctgt
          11761 aagggaaaca caaatggcca atcagatgct taatggtcat aggatgccag aaaatccaat
```

		11821	tacacctagt	cattgtattg	aaagggctgc	attgaaggaa	catttgaatc	atgttcctca
		11881	cgcaaaagcc	gcagtgatga	atggccaaat	gccccatagt	tacaggttgg	cgcaaaatcc
		11941	catcctacct	ccaaatcata	ttgaagggta	tcaagtgatg	gaaaatttga	gtgaacttgt
		12001	cacgacaaat	gactatctaa	ctgctagtcc	tttcagtcaa	actggagctg	caaataggca
	5	12061	gcataatatt	ggtgactcca	tgcatataca	tgcattggat	cctagaagag	agagtaatgc
		12121	ttcaagtggt	tcttggatat	cattaggtgt	gaactttaac	caacaaaata	atggatgggc
		12181	atctgcaggt	gctgccgatg	ctgcgagctc	acatgcccca	tatttttcag	aacctcacaa
		12241	aagaATGagg	acagcttatc	ttaacaatta	tccaaatgga	gtcgtgggac	atttttctac
		12301	ctcatctacg	gatttgtcaa	ataatgagaa	tgaaaatgtg	gcctcagcaa	tcaactcaaa
	10	12361	cgtttttacc	cttgctgatg	cacaaagatt	gatagcccgt	gagaaatcac	gagcttccca
		12421	aagaatgatc	agttttagat	catctaaaaa	tgatatggtt	aacagatcag	aaatggtcca
		12481	tcaacatggc	agacctgctc	cgcatggctc	tgcatgcagg	gagtctattg	aagtacctga
		12541	caaacagttc	gggctcatga	cagaagaact	cacacaatta	cctagtatgc	caaataaccc
		12601	acaaagggaa	aaatatattc	cgcaaactgg	aagttgccaa	cttcagtctt	tggaacatga
	15	12661	catggttaaa	gggcataact	t <u>g</u> gcaggtga	attgcataag	caagtaactt	cacctcaagt
		12721	tgttattcag	agcaatttct	gtgttacccc	tcctgatgtg	ctcggcagaa	gaaccagtgg
		12781	ggagcattta	agaaccctta	tagctccaac	acatgcatcg	acatgtaagg	acactctgaa
		12841	agctttaagt	tgtcaactgg	agagttctag	agacattatt	aggcctcctg	tcaatcctat
		12901	agggccatcc	tetgeegatg	ttccaagaac	tgataaccat	caagtcaagg	tttctgaaga
	20	12961	aaccgttaca	gccaaactcc	ctgagaagcg	aaaagtagga	cgtcccagaa	aagagttaaa
		13021	acctggtgag	aaaccaaaac	ctagaggccg	tccaaggaag	ggaaaagttg	ttggtggaga
		13081	acttgcatca	aaggatagtc	acactaatcc	attgcaaaat	gagagtactt	catgttctta
•		13141	tggtccttat	gcaggggagg	cttctgttgg	aagagcagtt	aaagcaaata	gagttggaga
		13201	aaacatttct	ggagctatgg	tatccctact	ggattcttta	gatattgtta	ttcaaaagat
	25	13261	aaaggtcttg	gacataaaca	aatcagaaga	ccctgtgaca	gctgaacctc	atggtgctct
	,	13321	tgtcccttac	aatggagaat	ttggtcctat	tgttcctttt	gagggaaag	tgaaaagaaa
		13381	acgctctcga	gccaaagtgg	atcttgaccc	tgtaactgct	ttaatgtgga	agttactaat
		13441	gggaccagat	atgagtgatt	gtgctgaa g g	tatggataag	gataaagaga	aatggctaaa
		13501	tgaagaaaga	aaaatattcc	aagggcgtgt	tgattcattt	attgctcgaa	tgcatctagt
	30	13561	tcaaggtatt	tctatcattt	taaaattgtt	ttcctaacat	gaacatgatg	gcttccatct
		13621	tgtgattgct	gccctcacat	tagtgaatgg	tctcaaatct	tcaatattta	ctgtgtaccc
		13681	aaatcctatt	tcttcatccc	aatatattca	tgtttgtact	cgtactgtcc	cattagactt
		13741	gcattgtgct	gtgaagatca	acacctttac	ttttaggatt	acctctatgt	ttgcaggaga
		13801	teggegtttt	tctccttgga	aaggatcagt	tgtagattct	gtagtgggag	tatttcttac
	35	13861	acagaatgtt	tcggaccatc	tttccaggtg	aataatgcct	agagcctatt	tgaaaactgt
		13921	gacttgactt	gcattgtgag	gttatgttgt	ttttctgtct	gactatttcc	tttttttca
		13981	gctctgcatt	tatggctctt	gctgcaaaat	ttcctgtaaa	gccagaagcc	tctgaaaaac
		14041	ccgcaaatgt	gatgtttcat	acaatttcag	aaaatggtga	ttgttctggg	ttgtttggta
	4.0	14101	attctgtcaa	gctacagggt	gagatccttg	ttcaggaggc	cagcaacaca	gcagcctctt
	40	14161	ttatcacaac	cgaggataag	gaaggaagta	acagtgtgga	attgcttgga	agttcttttg
		14221	gggatggagt	ggatggtgca	gcaggagttt	attctaatat	ttatgagaat	ctgccagcta
		14281	gactgcatgc	tactaggcgt	ccagtcgttc	aaactggaaa	cgctgtcgaa	gcggaagatg
		14341	ggtcactgga	gggtgttgtt	tcatcagaaa	actccactat	ttcatctcaa	aattcatcag
	4.5	14401	attatctatt	tcacatgtct	gatcatatgt	tttcgagcat	gttactaaat	ttcactgccg
	45	14461	aagacattgg	cagcagaaat	atgcccaaag	caacaagaac	cacatataca	gaacttctac
		14521	gaatgcagga	gctgaagaac	aagtctaatg	aaaccattga	atcatcagag	tatcatgggg
			ttccagtctc		_			
			aacatcagcc			_		_
	50		_			-		agagtgcttg
	50		_					gcctgtaaca
		14821	atgaaacaca	aaaggctgac	tctttaagca	acatgttata	tggtatagat	agatcagata

	14881	agactacttc	cctgtctgag	cctacaccaa	gaatcgataa	ctgttttcaa	ccattaagtt
	14941	cagagaaaat	gtcatttgct	agggaacagt	cctcttctga	aaattatctt	tcaaggaatg
	15001	aagctgaagc	tgcatttgtt	aaacagcatg	gaacatcaaa	tgtgcaaggt	gataatactg
	15061	tcaggacaga	gcaaaatgga	ggtgaaaatt	ctcaatcagg	atacagccaa	caggatgata
5	15121	atgttggatt	tcaaacagcg	acaaccagta	atctttattc	ttcaaactta	tgccaaaacc
	15181	agaaagcaaa	ttctgaagta	ctacacggag	tttcttccaa	cttgatagag	aattctaaag
	15241	atgacaaaaa	gacttcccc	aaagttccag	tcgatggatc	aaaagcaaag	aggccaagag
	15301	ttggggctgg	taaaaagaaa	acatatgatt	gggatatgtt	gagaaaagaa	gttctttaca
	15361	gtcatggtaa	taaagaaaga	tcccagaatg	ctaaggactc	aat t gattgg	gaaacaataa
10	15421	gacaagcaga	ggtgaaggaa	atatctgaca	caattagaga	gcgaggaatg	aataacatgc
	15481	tggcagaacg	gataaaagta	agtatggcat	aaaacagttt	acattgaaag	ttgacataac
	15541	tctagtcata	tgtgcatgca	tgctattcca	tatagatttg	cttatttgtt	ggaattccaa
	15601	gttttggatc	aaccatactc	atctttagca	attcatgttg	caggacttcc	taaaccgatt
	15661	ggtgagagac	catgggagca	tcgatcttga	gtggttgcgc	tatgtcgatt	cagataaagc
15	15721	gaagtaagct	aactaaattt	attttgagca	aacattcata	atgcaattgg	cccttgggca
	15781	ttctataatt	tgtcattttg	acctctgcat	tgcttagcaa	tgacaattgg	atgtagtgag
	15841	catgggtaat	aatgtaagca	atgacaattg	gatgtagtgg	gcatggttaa	taattgaaca
	15901	tgtctgtgtt	tgcgggataa	taatgcctat	cacctgtgag	cctgtgacat	gcaaaccttg
	15961	aacgttgaac	cttgaacccc	ctacctcgca	ctgtgtgctc	tcaaccaact	gagcaagtga
20	16021	gggaccttgt	tgtatggaaa	aaataatttt	aaataaccct	tgattcaacc	aaagcttcat
	16081	aaaagaatat	attttctatt	attcatttga	accagcggtt	gaaccagtga	accgatggtc
	16141	ttgctggtcc	ggatttaata	ataactatgg	ctagaacaga	ttagagcacc	gaatacttgc
	16201	gcgatgctaa	atatttcaat	ggggacacac	ctgctcgtgt	gttgcatcaa	ctacctaagc
	16261	cacacaggca	tggcaatcaa	atcagcttgc	ccatgtaaca	tcaactatct	gatcgcgaga
25	16321	aggccggagc	tctcacttga	tgtttgtcat	tcaaaaaata	gttattcacc	aatgcaatgt
	16381	caagctcccg	taaagaccat	gaatgtagtt	tatccttctt	tgatcaagtt	tttatttata
	16441	ttaaagtgtt	taccaatgta	atcctacatt	atttgtacct	ggtttttaca	tataaataca
	16501	ttgtaccttt	tgtgtttctt	ccagggacta	tctcttaagc	attagaggac	ttggacttaa
	16561	aagtgttgag	tgtgtgcgtc	ttttgacact	ccatcacatg	gcttttcctg	tatgtttcct
30	16621	ttcacaaata	attttcaaga	atcttcgttt	ctttatttct	ggagaagtgg	agattttatc
	16681	tgtatctgtt	gatgatgtag	gtggatacaa	atgttggtag	aatatgtgtg	aggcttggat
	16741	gggtgccact	tcagccccta	cccgagtctc	ttcagttgca	cctgttggag	atgtaagtat
	16801	cttaaatcca	ctggttggct	tcactaatgc	tggagagtga	taggagtttg	atcatctgct
	16861	attgaaggta	tccaatgctg	gagaacatac	agaaatacct	ctggccgagg	ttatgcaagc
35	16921	ttgatcaacg	gacattgtga	gttttagaaa	tgcagttaaa	aactatatat	ataagagcat
	16981	gtcattatct	gagagtgtaT	AA caggttct	tgatgatatg	taggtatgag	cttcactatc
	17041	aaatgataac	ttttggaaag	gtatgagaca	acaactttga	taaagtgaat	tcaacccaat
	17101	tactgtgttt	tgatggacca	tctgtgttac	tttccttcta	ggtattttgt	acaaaaagta
	17161	agcccaattg	caacgcatgc	ccaatgagag	ctgagtgcaa	gcactttgca	agtgcatttg
40	17221	ccaggtaatt	ctcaagatgt	acatattta	tatacattct	gtgaaatcac	ggtgatgatt
	17281	gttaggtatg	aacaattggc	tgagatcccc	ccctcccc	ctcccatcct	tttcctggtc
	17341	ctacaagttc	tcctaggcta	atttaactgg	tgcataccac	atttatgtta	ttttgataca
	17401	tcaaagatta	tgtttgtggt	tgtgaggcta	tattagtgtg	ttgtatgtaa	ctcagttttg
45	17461	caattgtagt	tttagttaga	acacgttgtt	ctctacattt	taataaatac	tttttgactg
	17521	gacatcaatg	actggtgtat	ttccgatata	aaaaggttga	ttgttgccga	gggatttcaa
	17581	ttcggtccga	ataggttcga	caaatgcagt	gggcctatta	gtttaagagt	gaaagttcta
	17641	tcagctgttt	gactccactg	tgacctttac	actttgtact	tttgaagaaa	cagactaacc
	17701	tgctcatatt	aaagtcttgg	aatgactcca	ttgcgacctt	tacgctttgt	attttagaag
	17761	aaacagacta	acctgttcat	attagagtct	tggaactgtg	tgtgtgtgtg	ttttttt
50	17821	ttttgggggg	gggggggcat	ggagatttaa	tccaacattc	ctggatgacc	ttatattggt
	17881	aatgatatgg	ttttttatg	atatagtgca	aggctcgctc	ttcctggacc	tgaagagaag

20941 tcctttttt tgccatttca gctgatagct actctccagt caaaatattt gccatctcta

40

```
21001 ttgaactttt cattgtcttc tgaatgtatc ttactcttgg atcattaata tttcattttg
          21061 tcacgatata gtggtatagg acaataaaat catgggaagt atttatttc atcaccaatc
          21121 tactcatata attttcaaat gacaattata aatatcttaa aaatatattg ttagttgtcc
          21181 tgtataaaat aattgtcaca ccctagtcca cagcgacaag aatttgtgtc tacaggctag
 5
          21241 agtgagtact ctagaagtat cttcatagga atcggaataa aatgccaatg tgaatgaaca
          21301 aggatatcaa gtataccctc aaaatctcta gagaggattg cgtaaatatg taggtgtaat
          21361 taaacaattg tttcatatgg agggttttct taaaggaggt acaagactta tcaatatggg
          21421 taaagtagtt tttatccata ggcattgttg gcagaaagct gcttagggta gaatgctact
          21481 ccctccgtcc cacaatataa gagattttga gtttttgctt gcaacgtttg accactcggc
10
          21541 ttattcaaaa atttttgaaa ttattattta ttttatttgt gacttacttt attattcaca
          21601 gtactttaag tacaactttt cgttttttat atttgcaaaa aaaattgtat aagacgagtg
          21661 gtcaaacgtt gtacgcaaaa actcaaaatc ccttatattg tgggacggag ggagtactta
          21721 tggatgcctt ttttgtccaa gatgtcagta acattttctt tcagggatgt ggatttttac
          21781 ttctttttc cctaactttt tcaggatttg tgtgcgtgag aggctttgat aggacatcaa
15
          21841 gagcacccag accactgtat gcaagactcc actttccagc aagcaaaatt accaggaata
          21901 aaaaatctgc aggttctgct ccaggaagag atgatgaata ggccatctgg aaaaccagaa
          21961 aggaaataaa gaggaggtac atatgatctg ccagaagatc actgacctga aatggatcgc
          22021 tgaccaataa gttgccgtag gcaattcaat tatttctggc catatacatc tgctgaaagt
          22081 tatgaactcc agccactgac gaattcgtgg tgctggtatt cttcggcaac atgatccatc
20
          22141 atacagatte tatgettggt tgttgeaage aattettatg eggtgacagt tgetgetgat
          22201 agggagaaaa ggcatgtccg gcggctcagc ggctctaact gtactttcat atgagtggaa
          22261 ccgattgttg tacatgtgaa aagtttgcca ttcaaaatgg tcattcatgt tgttaggtca
          22321 ttcatgtagt cgatgtcaaa ttaatcatca attatttgat ttgattcatt cacaagttta
```

25 SEQ ID NO:20

CORN(ZEA MAYS)DMT.1

>Corn DMT.1 660990 (688512 selclone ID);

EPDDPCPYLLSIWTPGETAOSIDAPKTFCDSGETGRLCGSSTCFSCNNIREMOAOKVRGT LLIPCRTAMRGSFPLNGTYFQVNEVFADHCSSQNPIDVPRSWIWDLPRRTVYFGTSVPTI FRGLTTEEIORCFWRGFVCVRGFDRTVRAPRPLYARLHFPVSKVVRGKKPGAARAEE

SEQ ID NO:21

>Corn DMT.1 cDNA 660990 (668512 selcone ID);

- 35 qaaccaqatqatccttqtccatatcttctttccatatqqaccccaqqtqaaactgcacaa tcgatcgatgccccaagac
 - attctgtgattcaggggagacgggtagactatgtggaagttcaacatgctttagttgcaa caatatacgagaaatgcagg
 - ctcaqaaaqtcaqaqqaacacttttgataccatqccqaacaqcaatqaqaqgaagcttcc
- cacttaatgggacgtatttt caagttaatgaggtatttgctgaccattgctcaagtcaaaatccaattgatgtcccacga agttggatttgggacctccc

aagacgaactgtttactttggaacctcagttcctacaatattcagaggtttaacgactga agagatacaacgatgctttt

ggagaggatttgtttgcgtgaggggctttgataggacagtgcgggcaccaaggccccttt atgcaaggttgcattttcct

5 gtcagcaaggttgttagaggcaaaaagcctggagcagcaagagcagaagaataatagaac attgaagaaatataggggtg

ctaaccagatgaggatggatagcccgaaatgagatgctgacccaataggtcgccaaatca cctccaaattctaacccaat

 $\tt gacttccatctgtaatgaatggcaataccttgaaaacctgtgatggagatgttttgtggc$

10 gacatgatctcttaaattag

attccgtctttggtaacagcctagctgttcttgttgagtcgcatattctttattctgaag atcaatatagcaaatggg

SEQ ID NO:22

15 CORN(ZEA MAYS)DMT.2

>Corn DMT.2 371537 (441428 selclone ID);

MITFGKVFCTKRQPNCNACPMRSECKHFASAFASARLALPAPQEESLVKLSNPFAFQNSS MHAMNSTHLPRLEGSIHSREFLPKNSEPIIEEPASPREERPPXTMENDIEDFYEDGEIPT IKLNMEAFAONLENCIKESNNELOSDDIAKALVAIXTEXASIPXPK

20

30

SEQ ID NO:23

>Corn DMT.2 cDNA 371537 (441428 selclone ID)

tatcagatgattacatttggaaaggtcttttgtaccaaaagacagccaaattgcaatgca tgcccaatgaggagtgagtg

25 caagcattttgcaagtgcatttgcaagtgcaaggcttgcacttcctgctccccaggagga aagcttagtgaagttgagca

agggagtttcttcctaagaactcagagccaataatcgaggagcctgcaagtccaagagag gaaagacctccakaaaccat .

ggaaaatgatattgaagatttttatgaagatggtgaaatcccaacaataaagcttaacat ggaagctttttgcacaaaact

tggagaattgcattaaagaaagcaataacgaactccagtctgatgatattgcaaaagcat tggttgctattarcactgaa rcagcttcsattcctgkaccgaaact

SEQ ID NO:24 5

Corn(Zea mays)DMT.3

>Corn DMT.3 218853;

MPRKPKRKAPASPARHDPSPEPYPSHASPCSAQCLVVRDALLAFHGFPEEFAAFRVLRLG GLSPNRDPRPSSPTVLDGLVTTLLSONTTDAISRRAFASLKAAFPSWDQVVDEEGKRLED AIRCGGLAATKAARIRSMLRDVRERRGKICLEYLRELSVDEVKKELSRFKGIGPKTVACV LMFYLQKDDFPVDTHVLRITKAMGWVPATASREKAYIHLNNKIPDDLKFDLNCLFVTHGK LCOSCTKKVGSDKRKSSNSACPLAGYCCIGEKLOOL

SEQ ID NO:25

WHEAT DMT.1 15

>Wheat DMT.1 614028 (887053 selclone ID);

MRAECKHFASAFASARLALPGPEEKSLVTSGNPIASGSCQQPYISSMRLNQLDWNANAHD HILDNROPIIEEPASPEPETAEMRESAIEDIFLDDPEEIPTIKLNFEEFAQNLKNYMQ VNNIEMEDADMSSALVAITPEAASIPTPRLKNVSRLRTEHOVYELPDSHPLLEGYDQREP

20 DDP

10

SEQ ID NO:26

>Wheat DMT.1 614028 (887053 selclone ID);

tgcccaatgagagctgaatgcaagcactttgcaagtgcatttgcaagtgctagacttgctcttcctggacctg 25 a agaga agagt tt g g tt acg t cag gaa acc ca att g ct t cag g g ag ct g c cag cag ca t a cat a ag t t ct at g c g t t t a a t ca a can a cat agaggagtttgcacagaatctcaagaattatatgcaagtcaataacattgaaatggaagatgctgatatgtcaagtgccttggttgccataactccggaagetgcatctatcccgactcctaggctcaagaatgttagtcgcctaagaacagagcatcaagtctatgaactgcc

30 ggactcacatccacttctggaaggatacgaccaaagagagcctgatgatccttg

SEQ ID NO:27

Wheat DMT.2

>Wheat DMT.2 568842 (908118 selclone ID);

NRVDESTVGGADKAASPKKTRTTRKKNTENFDWDKFRRQACADGHMKERKSERRDSVDWE AVRCADVQRISQAIRERGMNNVLSERIQEFLNRLVRDHGSIDLEWLRDIPPDSAKDYLLS IRGLGLKSVECVRLLTLHHLAFPVD

5 SEQ ID NO:28

>Wheat DMT.2 568842 (selclone ID 908118);

SEQ ID NO:29

15 WHEAT DMT.3

>Wheat DMT.3 611792 (838515 selclone ID);

NRKQVNEVFADHKSSYDPIYVAREQLWKLERRMVYFGTSVPSIFKGLTTEEIQQCFWKGF VCVRGFERETGAPRPLCOHLHVAASKVPRSRNAAAAGLNSDSAKASAP

20 SEQ ID NO:30

>Wheat DMT.3 611792(838515 selclone ID);

aatcgaaaacaagttaatgaggtatttgcagaacacaaatctagctacgatcccatatacgttgcaaggga gcagttatggaagttggaaagacgaatggtctactttggaacttcagtgccctccatattcaaaggtctaacaactgaagaaataca gcagtgcttctggaaaggatttgtctgtgtgcggggattcgagagggaaaccggggcaccaaggcctctatgccaacatctgca cgtcgcggctagcaaagtgccgagatcacgcaacgcggcagcagctgggctgaactcggattcagcaaaggcatcggctcca tgagtatcatcacaccggctatcgacctgtgcatgggtacgctagtgttggttcctgccgggcwacagccgttyttgtaggaaata aaccsctgcgcaaragaattatcatccagttggtytgagtgtatacttytgctgtagkaccttttttTaaaatccctgtgagctytattg taccttgaatttactttccgaccagtttatccgcttgcaaaraggcctttgttatgkaccggcatcttgttgtatatacatcatggttcctc traaaaacttgtcttgccakacgaccttacgt

30

25

SEQ ID NO:31

Wheat DMT.4

>Wheat DMT.4 615131 (861906 selclone ID);

15

MRSECRHFASAFASARLALPAPQEKSLVMSSNQFSFQSGGMPTPYSTVLPQLEGSAQGQD FCTNNSEPIIEEPASPAREECPETLENDIEDYDPDTGEIPLIKLNLQAFAQNLENCIKES NMDLGSDDIAKALVAVSTGSASIPV

5 SEQ ID NO:32

>Wheat DMT.4 615131 (861906 selclone ID);

tacttttggaaaggtgttctgtacaaaaaacaagccaaattgcaatgcttgtccaatgag aagcgaatgcaggcatttcgcaagtgccttcgcaagtgcacggcttgcacttcctgcacc tcaggagaaaagtttggtgatgtcgagcaatcaattcagtttccagagtggtggcatgcc aactccatactcaactgtgcttcctcagcttgagggaagtgcccagggacaggatttttg cactaacaattcagagccaattattgaggagccagcaagtccagcacgggaagaatgtcc agaaactcttgaaaatgata

ttgaagattacgatccagatactggtgaaatcccactaattaagcttaacttgcaagctt

ttgctcagaacttggaaaactgcattaaagaaagcaatatggatcttgggtctgatgata

tcgcgaaagcacttgttgctgttagcactggatcagcttcaattcctgtccc

SEQ ID NO:33

Soybean(Glycine max)DMT.1

>Soy DMT.1 449122 (557119 selclone ID);

MDSLDWDAVRCADVSEIAETIKERGMNNRLADRIKNFLNRLVEEHGSIDLEWLRDVPPDK
AKEYLLSIRGLGLKSVECVRLLTLHHLAFPVDTNVGRIAVRLGWVPLQPLPESLQLHLLE
LYPVLESIQKYLWPRLCKLDQETLYELHYQMITFGKXFCTKSKPNCNACPMRXECRHFAS
AFASARFALPGPEQKSIVSTTGNSVINQNPSEIISQLHLPPPENTAQEDEIQLTEVSRQL
ESKFEINICQPIIEEPRTPEPECLQESQTDIEDAFYEDSSEIPTINLNIEEFTLNLQN

25

30

20

SEQ ID NO:34

>Soy DMT.1 449122 (557119 selclone ID);

20

30

caaaaatatetetggeetegaetatgeaagetagateaggaaacaetatatgagetaeattaceagatgattaeatttggaaaggke ttetgtacaaaaageaaaceaaattgtaatgeatgeecaatgagaggagaat

SEQ ID NO:35

5 SOYBEAN(GLYCINE MAX)DMT.2

>Soy DMT.2 387990 (473695 selclone ID);

MRMTIDLVSQQSLTARLQLSILKDKLKIQCRKARGLDFGRNESSKIDSSPVKLRSREHGK
EKKNNFDWDSLRIQAEAKAGKREKTENTMDSLDWDAVRRADVSEIANAIKERGMNNMLAE
RIQSFLNLLVDKHGGIDLEWLRDVPPDQAKEFLLSIRGLGLKSVECVRLLTLHHLAFPVD
TNVGRIAVRLGWVPLQPLPESLQLHLLELYPVLESIQKYLWPRLCKLDQRTLYELHYQLI
TFGKVFCTKSK

SEQ ID NO:36

> Soy DMT.2 387990 (473695 selclone ID);

15 gaaaagataggatcattctcagatagcaactcagaaatagaagacctgtctagcgctgcc aagtacaatagttattataa

tagaatttctttcagtgagcttttagaaatggcaagttcaaccatgttgcatgaagttaa cagtcaaagaagcaaatcaa

ctgagaacttaggagatacatgtgatcagtctatagacatgaagcatgacaacctggcag aaaacttggaaaaatcggat


gttactcaaggctccgcagaagcacccatcaccaatggatatacttttaaaataacccca aactcaggagtacttgaggt

taactgttatgatcctctcaaaatagaagtcccatcaagtggctcctcaaagggtaaaga tgagaatgacaatagatcta

25 gtttcccaacagagtctgactgccaggctgcaattgtccattctcaaggacaaactgaag atccaatgcagGaaagcaag

agaaaaagaataactttgattgggatagtttaagaatacaagcagaagctaaggcaggga aaagagaaaagacagagaac

accatggactccttggactgggatgctgttagacgcgcagatgtcagtgaaattgccaat gcaatcaaagaaaggggcat



tgagagatgttccacctgatcaagcaaaagaattcttgctcagcataaggggattgggat
tgaaaagtgtggagtgtgta

5 cgactettaacactacaccatettgeettteeggtggacacaaatgttggacgtatagca gtaagattgggatgggtgee .

tctccagccactgccagagtcactacagttgcatcttctagaattgtacccagtgttgga gtccatacaaaaatatctct

ggccccggctctgcaagctagaccaaagaacattgtatgagctgcattaccagctgatta

10 catttggaaaggtcttctgt

actaaaagcaagcc

SEQ ID NO:37

SOYBEAN(GLYCINE MAX)DMT.3

15 >Soy DMT.3 657152 (546665 selclone ID);

INQAELQQTEVIRQLEAKSEINISQPIIEEPATPEPECSQVSENDIEDTFNEESCEIPTI KLDIEEFTLNLQNYMQENMELQEGEMSKALVALHPGAACIPTPKLKNVSRLRTEHYVYEL PDSHPLLNGWNKREPDDPGKYLLAIWTPGETABSIQPPESKCSSQEECGXLCNENECFSC NSFREAXFXDSXRDTPDTMSNSXXXGAFH

20

SEQ ID NO:38

>Soy DMT.3 657152 (546665 selclone ID);

tataaaccaagcagaacttcaacaaacagaagtgatcaggcaactagaagcaaaatctga aatcaacatcagccaaccta

25 ttattgaagagccagcaactccagagccagaatgctcccaagtatccgaaaatgatatag aggataccttcaatgaggaa

tcatgtgaaattcccaccatcaaactagacatagaagagttcactttgaacttacaaaac tatatgcaagaaaacatgga

30 tcctacacccaagctgaaga

atgtgagccggttgcgaacagagcattatgtttatgaactccctgattcacatccccttc tgaatgggtggaacaagcga .

gaacetgatgatecaggeaaataeettetagetatatggaeteeaggggagaeageagat tetataeageeaceagaaag

tcacagatagttcgagggacactcctgataccatgtcgaacagctwtgaragggag

5 SEQ ID NO:39

SOYBEAN(GLYCINE MAX)DMT.4

>Soy DMT.4 432980 (663678 selclone ID);

EAASIPMPKLKNVSRLRTEHCVYELPDTHPLLQGWDTREPDDPGKYLLAIWTPGETANSI OPPESKCSSQEECGOLCNENECFSCNSFREANSQIVRGTLLV

10

SEQ ID NO:40

>Soybean DMT.4 432980 (663678 selclone ID);

agaagctgcttccattcctatgcccaagctaaagaatgtgagccgattacgaacagagca ttgtgtttatgaactcccag

15 atacgcatcctcttctccaagggtgggacacacgagagcctgatgatccaggcaaatatc ttcttgctatatggactcca

ggtgagacagcaaattctatacagccaccagaaagcaaatgcagctctcaagaagaatgt ggccaactctgtaatgagaa

tga a tgtttctcgtgca a cagtttccgtga agca a attctcaga tagttagaggga cact

20 cctggtctgaatgcttatca

aaatcattgttttaaccatatgtagcttactaattcttatacattatgggaacaggggag ggaatacatctccatagaaa

ttcaagcattataatagactgacttgaatttatgataaatatgagcagataccatgt

>Medicago 6654943;

MELQEGEMSKALVALNQEASYIPTTKLKNVSRLRTEHSVYELPDSHPLLEGWEKREPDDP GKYLLAIWTPGETANSIQPPDRRCSAQDCGQLCNEEECFSCNSFREANSQIVRGTILIPC RTAMRGSFPLNGTYFQVNEVFADHESSLNPISVPRSLIWNLDRRTVHFGTSVTSIFKGLA TPEIQQCFWRGFVCVRSFERSTRAPRPLMARLHFPAS

SEQ ID NO:42

5

10

15

20

25

>Medicago 6654943 EST306265

GAGAACATGGAACTTCAAGAAGGTGAAATGTCAAAGGCCTTGGTTGCTCTAAATCAAGAA
GCTTCTTACATTCCTACAACGAAGCTGAAGAACGTGAGTCGGTTGCGCACAGAGCATTCT
GTTTATGAACTCCCAGATTCTCATCCTCTTCTGGAAGGGTGGGAAAAGCGAGAACCTGAT
GATCCAGGAAAATACCTTCTAGCTATATGGACGCCAGGTGAGACTGCAAATTCTATACAG
CCACCAGACAGAAGATGCAGCGCTCAAGATTGTGGCCAACTCTGTAATGAGGAGGAATGT
TTTTCGTGCAACAGCTTCCGTGAAGCAAATTCACAGATAGTTCGAGGGACAATCCTGATA
CCATGTCGAACAGCTATGAGAGGGAGCTTTCCGCTAAACGGAACCTATTTTCAAGTCAAT
GAGGTTTTTGCAGACCATGAATCAAGTCTTAATCCGATTAGCGTTCCCAGAAGTTTGATA
TGGAACCTTGATAGGAGGACAGTGCATTTTGGAACCTCCGTAACAAGCATATTCAAAGGT
TTAGCAACACCAGAAATTCAACAGTGCTTCTGGAGAGGGTTTGTCTGTGTGCGGAGCTTT
GAAAGGTCAACGAGAGCACCCCGTCCTTTAATGGCCAGACTTCCCAGCAAGC

SEO ID NO:43

>Tomato 12624037;

MELQEGEMSKALVALNQEASYIPTTKLKNVSRLRTEHSVYELPDSHPLLEGWEKREPDDP GKYLLAIWTPGETANSIQPPDRRCSAQDCGQLCNEEECFSCNSFREANSQIVRGTILIPC RTAMRGSFPLNGTYFQVNEVFADHESSLNPISVPRSLIWNLDRRTVHFGTSVTSIFKGLA TPEIQQCFWRGFVCVRSFERSTRAPRPLMA

SEQ ID NO:44

30 >Tomato 12624037 EST469495

10

15

25

30

TGAGGGATCATGGAAGTATTGACCTAGAATGGTTGAGAGATGTGGCCCCAGACAAAGCGA
AAGAGTATCTTTTGAGTATTCGTGGACTGGGTCTGAAAAGTGTAGAATGTGTGCGGCTAT
TAACACTTCATAACCTTGCTTTTCCAGTTGACACAAATGTTGGACGAATAGCTGTGAGAT
TAGGATGGGTTCCTCCCAACCACTTCCTGAGTCCCTGCAGTTGCATCTTCTTGAACTGT
ATCCAATTCTGGAGTCAATTCAGAAGTATCTCTGGCCACGACTCTGCAAGCTCGATCAGA
GAACACTGTATGAGTTGCACTACCACATGATTACCTTTGGAAAGGTTTTCTGCACCAAAA
GTAAGCCTAACTGTAATGCATGCCCACTGAGAGCTGAATGCAGACACTTTGCTAGTGCTT
ACGCAAGTGCAAGACTTGCCCTTCCTGGCCCAGAGGAGAAGAGTATAGTGAGTTCAGCAG
TTCCGATCCCTAGTGAGGGGAAATGCAGCCCATGCTATTACCCCCAG
AGCTGAAGTAGGGGATGCCGTACCCATATGCTCCAATTG

SEQ ID NO:45

>Barley 13256964;

MASETETFAFQAEINQLLSLIINTFYSNKEIFLRELISNASDALDKIRFESLTDKSKLDA QPELFIHIIPDKATNTLTLIDSGIGMTKSDLVNNLGTIARSGTKDFMEALAAGADVSMIG QFGVGFYSAYPCAERVXVTSKHNDDEQYGGEXQAGWLLYCGHVILLESPFGGVLRSPSTS RTNSWSTLERRAFKDLGKNTPSS

SEQ ID NO:46

-20 **>Barley 13256964** HVSMEI0014B12F

20

GTGGGTGGAAAATGGGTTCCTGGGGGGGCCCGGTTGAGGGATTGTTGGTCACATAAACA ACTATCGTCTTCTATCTTAGCACCTAATAGTCCTTCACATGAG

SEQ ID NO:47

5 >Corn 1BE511860;

LLEGFEQREPDDPCPYLLSIWTPGETAQSIDAPKTFCDSGETGRLCGSSTCFSCNNIREM QAQKVRGTLLIPCRTAMRGSFPLNGTYFQVNEVFADHCSSQNPIDVPRSWIWDLPRRTVY FGTSVPTIFRGLTTEEIQRCFWRGFVCVRGFDRTVRAPRALYAR

10 SEQ ID NO:48

>Corn 1BE511860 946063H01.Y1 946 -

SEO ID NO:49

>Cotton 11206330;

MQGNMELQEGDLSKALVALNPDAASIPTPKLKNVSRLRTEHYVYELPDKHPLLKQMEKRE

25 PDDPSPYLLAIWTPGETANSIQPPEQSCGSQEPGRLCNEKTCFACNSVREANTETVRGTI
LIPCRNAMRGSFSLNGT

SEQ ID NO:50

>Cotton 11206330 GA__EB0023J04F

30 CTCCGCCAGTGCATAACTTGCTTAAAGTAGGGCCTAATGTTGGCAACAATGAACCTATCA
TTGAGGAGCCTGCAACACCTGAACCAGAGCATGCAGAAGGATCAGAGAGTGATATTGAAG
ATGCAAGCTATGATGATCCAGATGAAATTCCCACAATAAAACTCAACATTGAAGAGTTCA
CAGCAAACCTACAGCATTACATGCAGGGCAATATGGAACTCCAAGAAGGGGACTTGTCAA
AGGCTTTAGTAGCTTTGAATCCTGATGCTGCTTCTATCCCTACTCCAAAATTGAAGAATG

10 SEQ ID NO:51

>Soybean 5606759

MGWVPLQPLPESLQLHLLELYPVLESIQKYLWPRLCKLDQETLYELHYQMITFGKVFCTK SKPNCNACPMRAECRHFASAFASARFALPGPEQKSIVSTTGNSVINQNPSEIISQLHLPP PENTAQEDEIQLTEVSRQLESKFEIYICQPIIEEPRTPEPECLQESXTDIEDAVYEDSS

15

20

25

SEO ID NO:52

>Soybean 5606759 SB95C12.

ACGAGCTTCCCGGTAGACACAAATGTCGGACGTATTGCCGTACGACTGGGATGGGTGCCT
CTGCAGCCACTGCCTGAGTCACTGCAGTTGCATCTCCTAGAATTGTACCCGGTGTTGGAG
TCAATACAAAAATATCTCTGGCCTCGACTGTGCAAGCTAGATCAGGAAACACTATATGAG
CTACATTACCAGATGATTACATTTGGAAAGGTCTTCTGTACAAAAAAGCAAACCAAATTGT
AATGCATGCCCAATGAGAGCAGAATGTAGACACTTTGCTAGTGCATTTGCAAGTGCAAGG
TTTGCACTGCCTGGACCAGAGCAGAAGAGTATAGTTAGCACAAACTGGAAATAGTGTGATT
AACCAGAACCCATCTGAAATCATCAGTCAGTTGCACTTGCCTCCACCTGAGAACACAGCC
CAAGAAGATGAAATTCAACTAACAGAAGTGAGCAGACAATTGGAATCAAAATTTGAAATA
TATATTTGCCAACCTATCATTGAAGAGCCCAGAACTCCAGAGCCAGAATGCTTGCAAGAA
TCACANACTGATATAGAGGATGCTGTCTATGAGGATTCAAGTG

SEQ ID NO:53

30 >Wheat 12019155

MFHCHGTRGSDLGFDLNKTPEQKAPQRRKHRPKVIKEAKPKSTRKPATQKTQMKENPHKK RKYVRKTAATPQTNVTEESVDSIVATKKSCRRALNFDLEHNKYASQSTISCQQEIDHRNE KAFNTTSDHKAKEPKNTDDNTLLLHEKQANNFQSE

>Wheat 12019155

AACAGTCAGGACAAAGGCAACAAGATCAGCAGTCAGGACAAGGGCAGCAACCGGGACAAA AACAAGGCAACAAGGTCAGCAGCCAGAACAAGGCCAGCAAGGTCAGCAACAAGACAAG GGGAGCAAGGTCAGCAGCCAGGACAAGGGCAACAAGGGCAGCAACCGGGACAAGGGCAGC CAGGGTACTACCCAACTTCTCCGCAGCAGTCAGGACAAGGGCAACCAGGGTACTACCCAA CTTCTCCACAGCAGTCAGGACAATTGCAACAACCAGCACAAGGGCAGCAACCAGGACAAG AGCAACAAGGTCAACAGCCAGGACAAGGGCAGCAACCGGGACAAGGGCAAGCCAGGGTAC TACCCAACTTCTCCGCAGCAGTCAGGACAAGAGCAACAGCTAGAACAATGGCAACAGTCA GGACAGGGGCAACCAGGGCACTACCCAACTTCTCCGTTGCAAGCCAGGACAAGGGCAACC AGGGTACTACCCAACTTCTCACAACAGATAGGACAAGGGCAGCAGCCAAGAACAATTTGC ACAACCAACACAAGGGCAACAANGGGCAGCAACCAAGGACAANGGGCAACAAGGTCAACA GCCCANGAAAAAGGCAACAAAGGTCAAGCAACCAAGNACAAGGGGCAGCAANCCAGGAC AAGGGCAGCCANGGTCCTACCCAACTTNTTTTGAGCAAGTCANGGAAAAGGGGCACCANC CNAGGANAAATGGGNACCACCCAGNACAAGGACAACCCCGGGTCTTCCCCAAANTTTTTN CN

20

25

10

15

SEQ ID NO:55

>Tomato 8106032

MSLAAHFPLKTDSTQKHEGNTGIIIEEPEECATDPNVSIRWYEDQPNQSTHCQDSSGVYN
TDSNEEKPAVNDSESSENSTECIKSAECSVILQSDSSREGSDLYHGSTVTSSQDRKELND
LPSSPSSVVSSEISAVIQASEGTDSSNFCSSTSFLKLLQMAGTSGAQGTRCTEHLHNQHK
GNXGQQPRTXGNKVNSPXKKATKVKQPXTRGSXPGQGQPXSYPTXFEQVXEKGHXPRXNG
XHPXOGOPRVFPKXF

SEQ ID NO:56

30 >Tomato 8106032 EST356474

25

AATACAGATTCAAATGAAGAAAAACCAGCTGTCAATGACTCTGAATCAAGTGAAAATAGC
ACAGAATGCATAAAATCAGCAGAATGTTCTGTAATTCTGCAATCAGATTCTTCTAGAGAA
GGCTCAGATCTGTATCATGGATCAACAGTTACAAGTTCCCAAGATCGAAAAGAGTTGAAT
GATTTGCCTTCTTCTCCGAGTTCTGTTGTTTCTTCTGAGATCTCTGCTGTTATTCAAGCT
TCAGAAGGAACTGACTCAAGCAACTTTTGCAGCTCCACTTCTTTTTTGAAGCTATTACAG
ATGGCAGGAACTTCAGGAGCACAAGGAACCAGGTGCACTGAACATCTAC

SEQ ID NO:57

5

>Corn 1AW042334;

10 DAHPLLQQLGLDQREHDDPTPYLLAIWTPDGIKEITKTPKPCCDPQMGGDLCNNEMCHNC
TAEKENQSRYVRGTILVPCRTAMRGSFPLNGTYFQVNEVFADHRSSHNPIHVEREMLWNL
ORRMVFFGTSVPTIFKGLRTEEIQQCFWRGFVCVRGFDMETRAPRPLCPHLHVIARPKA

SEQ ID NO:58

15 >Corn 1AW042334 614027C01.y1 614 -

SEQ ID NO:59

>Corn AW076298

PCRTAMRGSFPLNGTYFQVNEVFADHCSSQNPIDVPRSWIWDLPRRTVYFGTSVPTIFRG

30 LSTEQIQFCFWKGFVCVRGFEQKTRAPRPLMARLHFPASKLKNNKLTTEEIQQCFWRGFV

CVRGFDRTVRAPRPLYARLHFPASKVVRGK

>Corn AW076298 614065C03.Y1 614 -

15 SEQ ID NO:61

>Corn BE639158;

KNSEPIIEEPASPREERPPETMENDIEDFYEDGEIPTIKLNMEAFAQNLENCIKESNNEL QSDDIAKALVAISTEAASIPVPKLKNVLRLRTEHYVYELPDAHPLLQQLGLDQREHDDPT PYLLAIWTPDGIKEITKTPK

20

25

30

5

10

SEQ ID NO:62

>Corn BE639158 946021E09.Y1 946 -

TGAGCTGCATTATCAGATGATTACATTTGGAAAGGTCTTTTGTACCAAAAGACAGCCAAA
TTGCAATGCATGCTATGAATTCGACTCACCTACCTCGCCTTGAGGGGAGTATCCATTCAA
GGGAGTTTCTTCCTAAGAATTCAGÁGCCAATAATCGAGGAGCCTGCAAGTCCAAGAGAGG
AAAGACCTCCAGAAACCATGGAAAATGATATTGAAGATTTTTATGAAGATGGTGAAATCC
CAACAATAAAGCTTAACATGGAAGCTTTTGCACAAAACTTGGAGAATTGCATTAAAGAAA
GCAATAACGAACTCCAGTCTGATGATATTGCAAAAGCATTGGTTGCTATTAGCACTGAAG
CAGCTTCGATTCCTGTACCGAAACTAAAGAATGTGCTTAGGCTTCGAACAGAACACTATG
TGTATGAGCTTCCAGATGCACATCCACTTTTACAACAGCTAGGACTTGACCAACGGGAAC
ATGATGATCCTACCCCATACTTATTGGCCATATGGACACCAGATGGAATAAAGGAAATAA
CTAAGACACCAAAACCATGCT

>Corn T25243;

NHQPIIEEPLSPECETENIEAHEGAIEDFFCEESDEIPTINLNIEEFTQNLKDYMQANNV EIXYADMSKALVAITPDAASIPTPKLKNVNRLRTEHQVYELPDSHPLLEGFEQXEPDDPC PYLLSIWTPGELHNRSMP

SEQ ID NO:64

>Corn T25243;

SEO ID NO:65

>Corn AW453174;

20 FQGNEVFADHCSRQNPIDGPRSWIWDLPRRTGYFGTSGPTIFRGLTTEEIQRCFWRGFVC VRGFDRTVRAPRPLYARLHFPVSKVVRGK

SEQ ID NO:66

>Corn AW453174 660032D01.Y1 660 -;

>Corn BE509759;

NGTYFOVNEVFADHRSSHNPIHVEREMLWNLQRRMVFFGTSVPTIFKGLRTEEIQQCFWR GFVCVRGFDMETRAPRPLCPHLHIIARPKARKT

5

10

20

25

30

SEQ ID NO:68

>Corn BE509759 946021E09.X1 946 -

GGCTCTTTTTTAGGCAGGAAGTAATATGATTCCATGCATAGGTCGAGAGCTATTGATGTCATATCACAATAAACATGATG TTCATAAAACTGATATCTTTGCTGATTAGAGTACTTGCTCAGTTGCTGCTGTCTTGCGGGCCTTCGGCCTTGCTATAATG TGCAAATGGGGGCACAGAGGCCTTGGTGCTCTAGTCTCCATGTCGAATCCTCGCACACAGACAAATCCCCTCCAGAAGCA $\tt TTGTTGTATTTCTTGTTCTTAGACCTTTGAATATGGTGGGTACTGAAGTCCCGAAAAAGACCATGCGCCTTTGCAAGTCCTAGAGTCCCGAAAAAGACCATGCGCCTTTGCAAGTCCAAGTCCCGAAAAAGACCATGCGCCTTTGCAAGTCCAAGTCCCGAAAAAGACCATGCGCCTTTGCAAGTCCAAGTCCCGAAAAAGACCATGCGCCTTTGCAAGTCCAAGTCCCGAAAAAGACCATGCGCCTTTGCAAGTCCAAGTCCCGAAAAAGACCATGCGCCTTTGCAAGTCCAAGTCCCGAAAAAGACCATGCGCCTTTGCAAGTCCAAGTCCCGAAAAAGACCATGCGCCTTTGCAAGTCCAAGTCCCGAAAAAGACCATGCGCCTTTGCAAGTCCAAGTCCCGAAAAAGACCATGCGCCTTTGCAAGTCCAAGTCCAAGTCCCGAAAAAAGACCATGCGCCTTTGCAAGTCAAGTCCAAGTCCAAGTCCAAGTCAAGTCCAAGTCAAGTCAAGTCCAAGT$ TCCATAGCATCTCCCTTTCCACATGGATTGGGTTGTGGCTAGATCTGTGGTCAGCAAATACCTCATTGACTTGAAAGTAA **GTGCCATTAA**

15

SEQ ID NO:69

1AW017984; >Corn

VPRSWIWDLPRRTVYFGTSVPTIFRGLTTEEIQQCFWRGFVCVRGFDRTVRAPRPLYARL HFPASKVVRGK

SEO ID NO:70

1AW017984; >Corn

CCTGAAACAATCAAATAACGGCCGATGAGGTTACATTGTTTATAGTATATGATCAAAGAA CATGTATGACCATTGTACAAATAGGCCCATCTTCATGTTGTACAACAACTGATTCCACTC CAGAGTAAAGAATATGCGACTCACAAGAACATCTAAGCTGTTACCATAGATGGAACATAA TTTAACAGGTCATGTCGCCTCAAAAATTGGGTTAGAACTTGGAGGTGATTCGGCGACTTA TTGGGTCAGCATCTCATTTAGGGCTATCCATCCGTTTTGTTAGCTCCTCTATATTTCTTC GATGTACCTATTATTCTTCGACGCTTGCCGCTCCAGGCTTTTTGCCTCTAACAACCTTGC TGGCAGGAAAATGCAACCTTGCATAAAGGGGCCTTGGTGCCCTTACTGTCCTATCAAAGC CCCTCACAAACGAATCCTCTCCAAAAGCATTGTTGTATCTCTTCAGTCGTTAAACCTC TAAATATTGTAGGAACTGAGGTTCCAAAGTAAACAGTTCGTCTTGGGAGGTCCCATATCC AACTTCGTGGGAC